

SM PHARMACEUTICALS SDN. BHD.

FUNGI-STAT (ECONAZOLE NITRATE) PESSARY

DESCRIPTION:

A white colour, oval shaped pessary.

COMPOSITION:

Each pessary contains: Econazole Nitrate 150 mg

PHARMACODYNAMICS:

Fungistatic; may be fungicidal, depending on concentration; exact mechanism of action is unknown. Inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell wall membrane and altering its permeability; as a result, loss of essential intracellular elements may occur; also inhibits biosynthesis of triglycerides and phospholipids by fungi; in addition inhibits oxidative and peroxidative enzyme activity, resulting in intracellular buildup of toxic concentrations of hydrogen peroxide, which may contribute to deterioration of subcellular organelles and cellular necrosis. In *Candida albicans*, inhibits transformation of blastopores into invasive mycelial form.

Also has some activity against Gram-positive bacteria

PHARMACOKINETICS:

Absorption: Absorption is not significant when Econazole Nitrate is applied to the vagina.

Stratum corneum concentration: Far exceeded minimum inhibitory concentrations (MICs) for dermatophytes; inhibitory concentrations found in epidermis and as deep as middle region of dermis.

Elimination: Renal and fecal; <1% of applied dose recovered in urine and feces.

INDICATIONS:

Vaginitis caused by *Candida albicans* and other species of *Candida*.

CONTRAINDICATIONS:

Allergy to azoles.

SIDE EFFECTS / ADVERSE REACTIONS:

Vaginal burning, itching, discharge, or other irritation not present before therapy;
Hypersensitivity (skin rash or hives)

PRECAUTIONS / WARNINGS:

Pessary should only be inserted into the vagina and should not be eaten or swallowed.
Avoid contact with contraceptive diaphragms and condoms since pessary may damage the rubber.

Do not use if this medicine if:

- Patients have experienced an allergic reaction (hypersensitivity) to Econazole medicine or any medicine used to treat thrush or other fungal infections.
- Patients believe or suspect symptoms are not due to Candida (Thrush) but another sexually transmitted disease. Please consult a doctor.
- Patients should not use this medicine without first consulting a doctor if they have: abnormal or irregular vaginal bleeding, blood stained discharge, vulval/vaginal sores or blisters and tummy pain or pain on passing urine.

Patients under 16 years old or over 60 years should not use this medicine without consulting the doctor.

Pregnancy

Patients should tell the doctor if she is pregnant or may be pregnant.

Econazole pessaries are effective in the treatment of vaginal candidosis in pregnant woman. However, safety of the drug in pregnancy has not been established and it should not be used unless the expected benefit outweighs any potential risk. If there is a strong need for the use of Econazole pessaries during pregnancy, in particular during the first trimester, the patient should be advised to exercise caution in the insertion of a vaginal applicator.

Lactation

It is not known whether econazole is distributed into human breast milk. However, problems in humans have not been documented.

DRUG INTERACTIONS:

There are no known drug interactions or physical incompatibilities between Econazole Pessary and other drugs, but you should tell your doctor if you are using, or have recently used, anything else to treat your present symptoms.

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION:

150 mg (one pessary) once a day at bedtime for 3 consecutive days. May be repeated if needed. It is important to use this medicine at the right times and in the right way. Insert the pessary into the vagina.

If sensitisation or irritation occurs, treatment with vaginal azoles should be discontinued

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTES:

No cases of deliberate overdosage have been reported. If one or more pessaries are eaten or swallowed, gastric emptying is recommended by giving Ipecac syrup to induce vomiting. This can be followed by giving activated charcoal as an absorbent to the drug.

DOSAGE FORMS AND STRENGTHS AVAILABLE:

Fungi-Stat Cream 1% w/w; Fungi-Stat Pessary 150 mg

PACKING / PACK SIZES:

Strip pack of 3x10 pessaries /box.

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS:

Store below 30°C. For vaginal use only. Continue medicine for full time of treatment.

SHELF LIFE: 3 years

NAME AND ADDRESS OF MANUFACTURER:

SM PHARMACEUTICALS SDN BHD (218620-M)
Lot 88, Sungai Petani Industrial Estate
08000 Sungai Petani, Kedah Darul Aman, Malaysia.

24.01.2020