



消渴靈片

Xiao He Ling Pian

功能：传统用于 益氣，利尿，消渴。

服法：成人：每次 4-7 片，每日 3 次，食后。

Each tablet (445.0mg) mengandungi:

Rhizoma atractylodis chinensis	10.68 mg
Radix glycyrrhizae glabra	23.48 mg
Radix salviae miltiorrhizae	21.33 mg
Radix achyranthis bidentatae	6.40 mg
Ramulus loranthis mulberry mistletoe	10.68 mg
Radix paeonia alba	8.53 mg
Rhizoma ligustici chuanxiong	8.53 mg
Fructus schizandrae chinensis	19.21 mg
Poria cocos wolf	10.68 mg
Radix astragali seu hedyari	55.49 mg
Fructus corni officinalis	10.68 mg
Radix atractylodis macrophalae	8.53 mg
Herba taraxacum mongolicum hand mazz	19.21 mg
Rhizoma polygonatum sibiricum redoute	12.81 mg
Rhizoma coptidis chinensis	8.53 mg
Rhizoma dioscoreae opposita thunb	10.68 mg
Radix rehmanniae ghunosa	12.81 mg
Rhizoma alismatis orientalis	8.53 mg
Radix scrophulariae ningpoensis	10.68 mg
Radix rehmanniae praeparata	12.81 mg
Radix angelica sinensis	10.68 mg
Radix scutellariae baicalensis	12.81 mg
Semen coix lachryma jobi L.	10.68 mg
Radix bupleuri chinensis	8.53 mg
Cortex moutan radidis	8.53 mg
Fructus lycii barbarum	10.68 mg
Radix ophiopogonis japonicus	8.96 mg
Radix puerariae lobata	8.96 mg
Gypsum fibrosum	10.68 mg
Herba epimedium brevicornum maxim	10.68 mg
Herba dendrobium nobile lindl	8.53 mg

Description: Brown, round biconvex tablet. Dm=10mm

Indication: Traditionally used for invigorating vital energy, improve urination and sweating.

Digunakan secara tradisional menambah tenaga, melawaskan kencing dan peluh.

Adult: 4-7 tablets each time, three times daily, after food.

Dewasa: 4-7 tablet setiap kali, tiga kali sehari, selepas makan.

This is a traditional preparation. Please consult your doctor/pharmacist before taking this product.

Ini ubat tradisional. Sila hubungi doktor/ahli farmasi anda sebelum mengambil produk ini.

傳統草藥配方，服用本产品前请咨询医生/药剂师。

Keep below 30°C. Protect from light and moisture.

Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Simpan di bawah 30°C. Lindungi daripada cahaya dan kelembapan.

收藏於干燥 30°C 以下。避免光照和潮湿。

放在儿童不能接触的地方。

WARNING: Not to be taken by babies, children under 12 years of age, pregnant women or lactating mothers. Consult your practitioner if you have conditions such as:

glucose-6-phosphate dehydrogenase (G6PD) deficiency, haemolytic anemia, glaucoma, diabetes, high blood pressure, history of cardiovascular disease and if you are using paclitaxel, cyclosporin, or other chemotherapeutic agents.

Packing: Per bottle of 120's, 240's and 1000's



Manufactured by product registration holder:

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong perusahaan Utama 1,
Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang.

Regd. no : MAL20040585T

Date of revision: 20/06/2019