



OXYMETAZOLINE 0.025% NASAL DROPS

COMPOSITION:

Contains Oxymetazoline HCl 0.025% w/v
(Benzalkonium Cl 0.01% w/v as preservative)

PRESENTATION:

Clear, colourless liquid.

INDICATIONS:

Relief of nasal congestion in acute colds, paranasal sinusitis and otitis media.

PHARMACOLOGY:

Mechanism of Action:

Oxymetazoline is a direct-acting sympathomimetic agent which has a vasoconstrictor effect on mucosal blood vessels. Sympathomimetic agent mimic the actions produced by stimulation of postganglionic sympathetic or adrenergic nerves, including vasoconstriction of blood vessels supplying skin and mucous membranes. Topical use of Oxymetazoline has the advantage of selective of site of action and more rapid action than do oral agents. It acts within a few minutes and the effects last for up to 12 hours. Nasal congestion is frequently a symptom of conditions such as rhinitis. Sympathomimetic agents are widely employed as nasal decongestants to provide symptomatic relief. They act by vasoconstriction and redistributing local blood flow to reduce oedema of the nasal mucosa thus improving ventilation, drainage and nasal stuffiness.

Pharmacokinetics:

Most sympathomimetic agents undergo enzymatic degradation in the gut and first-pass metabolism of the liver. They are inactivated by the enzymes catechol-O-methyltransferase (COMT) and monoamine oxidase (MAO). The metabolites are excreted in the urine mainly as their glucuronide and ethereal sulphate conjugates. Oxymetazoline, however, is an imidazoline derivative and is more resistant to the inactivating enzymes of liver and other tissues. It is thus longer action and metabolised at a slower rate than the catecholamines such as ephedrine or phenylephrine. When used in the form of nasal drops it acts with a few minutes and the effects lasts for several hours.

DIRECTION FOR USE:

For nasal use.

Axcel Oxymetazoline 0.025% Nasal Drops may only be administered to children aged between one and six years and should not be used in infants under 1 year of age. Unless otherwise prescribed, instill 1-2 drops of Axcel Oxymetazoline 0.025% Nasal Drops into each nostril 2-3 times daily. Unless specifically prescribed by the doctor, Axcel Oxymetazoline 0.025% Nasal Drops should only be used for short periods of time (up to 10 days). A treatment-free period of several days should precede any repeated use. Permanent use of decongestant rhinological agents may attenuate their effect. The abuse of local rhinological agents may cause mucosal atrophy and reactive hyperaemia with rhinitis medicamentosa. Longer use of oxymetazoline may cause damage to the mucosal epithelium with inhibition of ciliary activity. This may possibly result in

irreversible damage to the mucosa with rhinitis sicca. Long term use and overdosage must be avoided, especially in children. Medical supervision is indicated in patients with chronic rhinitis. Dosage higher than recommended may only be used under supervision.

CONTRAINDICATION:

Contraindicated in patients with known hypersensitivity to Oxymetazoline & Benzalkonium Chloride. Should not be used where there is inflammation of the skin and mucosal of the nasal vestibule with incrustation (rhinitis sicca).

PRECAUTION:

May cause hypertensive crisis if used during treatment with a monoamine-oxidase inhibitors (MAOI's) and it should not be given to patients receiving such treatment or within 14 days of its termination. Prolonged use of nasal decongestant drops may cause rebound congestion and drug-induced rhinitis. Long-term use and overdosage must be avoided, especially in children. Dosage higher than recommended may only be used under medical supervision. This product should not be used by patients who have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland without careful clinical consideration.

Pregnancy and Lactation:

Safe use of oxymetazoline hydrochloride has not been established with respect to adverse effects upon fetal development. Therefore it is not recommended for use in pregnancy.

Use in infants:

Should not be used in children younger than 1 year of age.

SIDE EFFECTS:

Oxymetazoline may occasionally cause local stinging or burning, sneezing and dryness of the mouth and throat. Prolonged or excessive use may cause rebound congestion and drug induced rhinitis.

DRUG INTERACTIONS:

Sympathomimetic may increase blood pressure and therefore special care advisable in patient receiving anti-hypertensive therapy. May cause hypertensive crisis if used during treatment with monoamine-oxidase inhibitor.

OVERDOSAGE AND TREATMENT:

Overdosage may cause CNS depression, leading to coma and marked reduction in body temperature, especially in infants. Systemic effects such as tachycardia, somnolence and shock-like states have been reported in infants and children after administration of a large number of drops. In case of overdosage, the appropriate symptomatic and supportive treatment should be given.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

PACK QUANTITIES:

Available in 10ml amber glass bottle.

Further information can be obtained from doctor, pharmacist or the manufacturer.

Product Registration Holder & Manufactured By:

 **Kotra Pharma (M) Sdn. Bhd.**
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

PMSPAAXC08100 - 050924(05)

Date of revision : 01/03/24

MAL20040821AZ