

GLUCOSAMINE CAPSULE 250MG

Ingredient(s):

Each capsule contains:
Glucosamine Sulphate 250mg

Pharmacology (Summary of Pharmacokinetics and Pharmacodynamics):

1. Glucosamine Sulphate stimulates the anabolic metabolism of osteo-cartilaginous tissues through stimulation of the biosynthesis of the mucopolysaccharides (which are the essential components of the cartilage ground substance). It provides lubricant activity through improvement of synovial fluid viscosity and increase of synovial fluid production.
2. Glucosamine possesses arthrotrophic metabolic activity by stimulating the anabolic metabolism of osteo-cartilaginous tissues and improvement of lubricant activity. The pharmacokinetics of glucosamine sulphate was investigated in 6 healthy male volunteers (2 per administration route) using ¹⁴C uniformly labelled glucosamine sulphate and administering it in single dose by intravenous (i. v.), intramuscular (i. m.) or oral route. After oral administration a proportion close to 90% of glucosamine sulphate is absorbed. Free glucosamine is not detectable in plasma. The radioactivity incorporated in the plasma proteins follows pharmacokinetics patterns which are similar to those after i. v. or i. m. administration, but its concentration in plasma is about 5 times smaller than after parental administration. The AUC after oral administration is 26% of that after i. v., or i. m. administration. The plasma levels of radioactivity investigations in rats and dogs showing that also in man glucosamine sulphate is prodrug for glucosamine that is well absorbed after oral administration and that diffuses into several tissues, including bones and articular cartilages.

Indication(s):

As adjuvant therapy for osteoarthritis.

Dosage and Administration:

2 capsules 3 times a day for 6 to 8 weeks. Maintenance: 2 capsules 2 times a day for 3 to 4 months.

Contraindication(s):

Patients who are hypersensitive to glucosamine.

Precaution(s) / Warning(s):

In pregnancy and lactation, Glucosamine should be given under medical supervision.

Side Effect(s) / Adverse Reaction(s):

The most frequently observed adverse events occurring after glucosamine treatment were mild digestive problems, including epigastric tenderness, heartburn, diarrhea and nausea.

CARDIOVASCULAR:

Peripheral Oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

CENTRAL NERVOUS SYSTEM:

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

GASTROINTESTINAL:

Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

SKIN:

Skin reaction such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Symptoms and Treatment for Overdosage, and Antidote(s):

None known.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A #1 red cap and white body hard gelatin capsule, both sides are impressed with " YSP " containing white to slightly brownish powder.

Plastic bottle of 30's, 80's and 100's.

Plastic bottle of 500's and 1,000's (for export only)

DERIVED FROM SEAFOOD

Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 13 Nov 2024

Item Code	: Draft 13 Nov 2024
PM Code	: Ins.P- GSC
Measurement(mm) ± 2mm:	260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: mm
Colour of Artwork	:  K
Source of Artwork	: MAC 3 / CC

L
260mm
(±2mm)

W
110mm (±2mm)