

1 page

2 page

leaflet as booklet

3 page

4 page

5 page

6 page

LfB3234v1Gal200s1-MAL0320 PACKAGE LEAFLET FOR: Gallifen 200 mg/ml suspension for use in drinking water for chickens 1. PHARMACODYNAMICS Fenbendazole is an anthelmintic belonging to the benzimidazole-carbamate group. It acts by interfering with the energy metabolism of the nematode. Fenbendazole inhibits the polymerisation of tubulin to microtubules. This interferes with essential structural and functional properties of the cells of helminths, such as formation of the cytoskeleton, formation of the mitotic spindle and the uptake and intracellular transport of nutrients and metabolic products. Fenbendazole is active and has a dose activity on adult <i>Heterakis gallinarum</i> and <i>Ascaridia galli</i>. 2. PHARMACOKINETICS After oral administration fenbendazole is only partially absorbed. Following absorption, fenbendazole is rapidly metabolised in the liver mainly to its sulphoxide (oxfendazole) and further to its sulphone (oxfendazole sulphone). In chickens oxfendazole sulfone is the main component detected in plasma, accounting for about 3/4 of the total AUC (i.e. the sum of the AUC for fenbendazole, oxfendazole and oxfendazole sulphone). Fenbendazole and its metabolites are distributed throughout the body, reaching highest concentrations in the liver. 3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENT Suspension for use in drinking water	Each ml of white to almost white suspension contains : Active substance: Fenbendazole 200 mg Excipients: Sodium benzoate 3 mg 4. INDICATION(S) Treatment of chickens infected with <i>Heterakis gallinarum</i> (adult stages) or <i>Ascaridia galli</i> (adult stages). 5. CONTRAINDICATIONS Do not use in known cases of hypersensitivity to the active substance or to any of the excipients 6. ADVERSE REACTIONS None known. 7. TARGET SPECIES Chickens. 8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION For use in drinking water. Shake well before use. The dose is 1.0 mg fenbendazole per kg body weight per day (equivalent to 0.005 ml Gallifen suspension). This dose has to be administered on 5 consecutive days. Dose calculation: The required daily amount of product is calculated from the total estimated body weight (kg) of the entire group of chickens to be treated. Please use the following formula: ml product/day = total estimated body weight (kg) of chickens to be treated x 0.005 ml 9. ADVICE ON CORRECT ADMINISTRATION To ensure administration of the correct dose, body weight should be determined as accurately as possible. Before allowing animals to have access to the medicated water, the water delivery system should be drained, if possible, and flushed with	the medicated water to ensure accuracy of dosing. This procedure may need to be performed on all treatment days. The uptake of medicated water depends on age and clinical conditions of the chicken, ambient temperature and light regime. In order to obtain the correct dosage concentration of the product should be adjusted accordingly. Follow the instructions described below to prepare the medicated water. Use a sufficiently accurate commercially available measuring device. For each treatment day the medicated water needs to be freshly prepared. For use in medication tank: Add the calculated amount of product to 40 to 80 % of the chicken daily water ration. Stir until content in the medication tank is visibly homogeneous. The medicated water appears hazy. No further stirring during administration is necessary. For use in dosing pump: Add the calculated amount of product to the unmedicated water in the stock suspension container of the dosing pump. The volume of unmedicated water in the stock suspension container has to be calculated taking as a basis the preset injection rate of the dosing pump and 40 to 80 % of the chicken daily water ration. Stir until content in the stock suspension container is visibly homogeneous. The medicated water appears hazy. During treatment all animals must have solely but unrestricted access to the medicated water. During treatment, after complete consumption of the medicated water, animals must be allowed access to un-medicated drinking water as soon as possible.	Ensure that the total amount of medicated water offered is consumed. 10. WITHDRAWAL PERIOD Meat and offal: 6 days Eggs: zero days 11. STORAGE Store below 30°C (room temperature). Store in the original container protected from direct sunlight. Do not freeze. Protect from frost. 12. SHELF-LIFE Shelf life of the veterinary medicinal product as packaged for sale: 36 months. Shelf life after first opening the immediate packaging: 6 months when stored below 30°C (room temperature). Shelf life of the medicated drinking water: 24 hours. 13. WARNINGS AND PRECAUTIONS Special warnings for each target species Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy: - Too frequent and repeated use of anthelmintics from the same class, over an extended period of time. - Under dosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any). Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.	Special precautions for use <u>Special precautions for use in animals</u> The safety of the product at overdose has not been evaluated in chickens less than 8 weeks old. The efficacy of the veterinary medicinal product at the recommended dosage is not sufficient for the treatment of infections with <i>Capillaria</i> spp. The absence of <i>Capillaria</i> spp infestation should be confirmed prior to use of the product. In case of <i>Capillaria</i> infestation another appropriate anthelmintic veterinary medicinal product should be used. <u>Special precautions to be taken by the person administering the veterinary medicinal product to animals</u> Embryotoxic effects cannot be excluded. Pregnant women must take extra precautions when handling this veterinary medicinal product. This veterinary medicinal product may be toxic to humans after ingestion. This product may cause eye irritation. Contact with the skin and the eyes or accidental ingestion of the product should be avoided. Do not smoke, eat or drink when handling the veterinary medicinal product. In the event of accidental ingestion, rinse mouth with plenty of clean water and seek medical advice. In the event of accidental contact with the skin or eyes, rinse with plenty of clean water and seek medical advice. Wash hands after use. Use during pregnancy, lactation or lay Can be used during lay. Interaction with other medicinal products and other forms of interaction None known. Overdose (symptoms, emergency procedures, antidotes), if necessary No adverse reactions have been	observed at up to 5-fold overdose in broilers (aged approximately 8 weeks). No adverse reactions have been observed at up to 3-fold overdose in layers and breeders. Incompatibilities In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products. 14. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. The product should not enter water courses as this may be dangerous for aquatic organisms. 15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED February 2020. 16. PACKAGING AVAILABLE Bottles of 1 litre; Canisters of 2.5 litres and 5 litres. Not all pack sizes may be marketed. NAME AND ADDRESS OF THE MANUFACTURER Biovet Joint Stock Company 39 Petar Rakov Street, 4550 Peshtera, Bulgaria NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER Pahang Pharmacy Sdn. Bhd. (10826-K) Lot 5979 Jalal Teratai, 5 ½ Mile Off Jalan Meru, 41050 Klang, Selangor Darul Ehsan, Malaysia
---	---	--	---	--	---