

ROBENXIN 66G/KG POWDER

Description and Composition

ROBENXIN 66G/KG POWDER is an off white to yellowish powder. Each kg contains 66g of Robenidine Hydrochloride as its active ingredient.

Pharmacodynamics

Antiprotozoal containing robenidine hydrochloride is used for anticoccidial treatment in broiler production. Shows anticoccidial activities against: *Eimeria acervulina*, *E. maxima*, *E. tenella*, *E. necatrix*, *E. brunetti* and *E. mivati*.

Acting robenidine hydrochloride is the result of dual action which is shown at different stages of the parasite's development in the intestinal mucosal part of broiler chickens. The first mechanism of action is reflected like this: robenidine hydrochloride inhibits the development of the first generation schizont, and the other is reflected by killing another generation schizont and possibly merozoite.

Also, robenidine hydrochloride acts on gametogenesis parasites. Dual mechanism of action is particularly advantageous for anticoccidial drugs, because the risk of relapse is reduced in case of bird's stress and following feed termination. Combination of coccidiocides and coccidiostatic effects is also important factor in minimizing the possibility of clinical manifestation of coccidiosis after the discontinuation of robenidine hydrochloride.

By using histochemical techniques, it has been confirmed that robenidine hydrochloride has a negative effect on the development of the first generation of schizont, first generation of merozoite and the second generation of schizont. It was also confirmed that robenidine hydrochloride gives effect in preventing anticoccidial protein synthesis.

Pharmacokinetics

Reabsorption

After oral application, robenidine is partially absorbed from the gastrointestinal tract (20 to 40%) and rapidly excreted, mainly it is not metabolized.

Distribution

After a single dose (5 to 8 times higher than the recommended dose), robenidine distribution in tissues and the medicine reaches the highest concentration for 8 hours in the muscles, liver, kidneys and blood, fat (as in subcutaneous) and bile during 24 hours after application.

Metabolism

Biotransformation of robenidine mostly occurs in the liver. Emerging metabolites are not active.

Elimination

Unabsorbed robenidine is excreted in faeces and emerging metabolites.

Indication

As an aid in the prevention of coccidiosis caused by *E. mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima* and *E. necatrix* in broiler chickens.

Recommended Dose

30 grams of Robenidine Hydrochloride (0.0033 percent) per ton (907kg) of feed. Feed continuously as sole ration.

Route of Administration

For oral administration via feed.

Contraindication

None known.

Warning and Precautions**Warnings:**

Do not use in birds which are producing or may in the future produce eggs or egg products for human consumption.

Special precautions to be taken by the person administering the drug to the animals:

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale dust. If product on skin, wash area immediately with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, face and arms thoroughly with soap and water. Wash hands after use. When opening container and mixing (into stock feed) wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles and disposable dust mask. After each day's use, wash gloves, goggles and contaminated clothing.

Interaction with Other Medicaments

Do not use concurrently with other anticoccidial drugs.

Pregnancy and Lactation

Not applicable.

Side Effects

If the medicine is applied in accordance with the instructions, side effects were not recorded.

Symptoms and Treatment of Overdose

The highest tested dose of 300 mg/kg feed (8 times the maximum authorised level), resulted in significantly depressed weight gain and increased feed conversion rate compared to untreated birds. No other adverse effects were observed. Gross pathology and histopathological investigations (liver, kidney, spleen, gonads, brain and muscles examined) did not show any substance-related alterations.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Period

Chicken (broilers): 5 days

Do not feed to chickens producing eggs for food.

Packing

10kg and 25kg.

Revision date: 15.1.2020

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com