

DURACIN 10G/KG POWDER

Description and Composition

DURACIN 10G/KG POWDER is a brown powder. Each kg contains 10g of Maduramicin ammonium as its active ingredient.

Pharmacodynamics

Maduramicin is a polyether antibiotic that belongs to the lipophilic class of antibiotics known as ionophores. These compounds form electrically neutral complexes with monovalent and divalent cations. The lipophilic character of these antibiotics render them soluble in lipid components and they catalyze electrically silent exchanges of cations for protons or other cations across a variety of biological membranes. The ionophorous nature of these antibiotics results in a wide range of biological activities

Pharmacokinetics

Maduramicin was excreted rapidly in chickens. Steady state was observed after three days in the excreta, but after six days in plasma. Maduramicin was the main compound excreted (26%). O-demethylation represented the main metabolic pathway of maduramicin common to chickens.

Indication

For the prevention of coccidiosis in broiler chickens caused by *Eimeria acervulina*, *E. tenella*, *E. brunetti*, *E. maxima*, *E. necatrix* and *E. mivati*.

Recommended Dose

454g to 545g of Duracin 10g/kg Powder which is equivalent to 4.54g to 5.45g (5ppm to 6ppm) Maduramicin ammonium per ton of feed, per day for 5 consecutive days orally in feed.

Route of Administration

For oral administration via feed.

Contraindication

Dangerous for equine species. This feeding stuff contains an ionophore: simultaneous use with certain medicinal substances (e.g. tiamulin) can be contraindicated.

Warning and Precautions

Special warning for target species:

1. Treated chickens must not be slaughtered for use in food for at least five days after the latest treatment with this drug.
2. Do not feed to replacement, laying or breeding chickens.

Special precautions to be taken by the person administering the drug to the animals:

Handle with care. When mixing feed, operators should wear protective clothing, safety glasses and face masks. Wash hands thoroughly after feed preparation.

Other Precautions:

1. Use only as recommended. Do not exceed the recommended level of 5ppm (mg/kg) as reduced weight gains may result with overdosage.
2. Do not use this feed for treatment of outbreaks of coccidiosis.

Interaction with Other Medicaments

This feeding stuff contains an ionophore: simultaneous use with certain medicinal substances (e.g. tiamulin) can be contraindicated.

Clinically important interactions between ionophore anticoccidials and the antibiotic tiamulin (and other antibiotics) are well-known phenomena in chickens. The severity of adverse effects varies depending on the ionophore itself, reaching from increased mortality to only transient performance depression.

Pregnancy and Lactation

Studies designed to assess the mutagenic effects of maduramicin produced negative results. Studies designed to assess reproductive effects including teratogenicity demonstrated a lack of any adverse effects on reproductive performance and a lack of teratogenicity at any dose level of maduramicin tested.

Side Effects

No adverse commercial effects on litter conditions, disease outbreaks or feathering were observed in any of the pivotal studies of floor pen safety study in broiler chickens.

Symptoms and Treatment of Overdose

Doses greater than 6ppm showed a statistically significant decrease in weight gain as compared to the non-medicated control group.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Period

5 days.

Packing

10kg and 25kg.

Revision date: 14.1.2020

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com