

ESPACOX 50 mg/ml

Oral suspension
Toltrazuril

Composition

1 ml contains: **Active substance:** Toltrazuril, 50 mg. **Excipients:** Sodium benzoate, 2.1 mg and Sodium propionate, 2.1 mg (as preservatives); Excipients, q.s.
White or yellowish oral suspension.

Indications

Cattle: Prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Pigs: Prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: Prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria cran-dallis* and *Eimeria ovinoidalis*.

Contraindications

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

Adverse reactions

None known. If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

Target species

Cattle (calves: dairy calves, beef calves, fattening calves), pigs (piglets, 3-5 days old) and sheep (lambs).

Dosage for each species, route and method of administration

Oral use.

All species: To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Cattle: Each animal should be treated with a single dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

Pigs: Each pig should be treated on day 3 - 5 of life with a single dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended.

Sheep: Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

Shake well before use.

Withdrawal period

Meat: Cattle: 63 days; Pigs: 77 days; Sheep: 42 days.

Milk: Not authorized for use in animals producing milk for human consumption.

Storage precautions

Keep out of the sight and reach of children.

Do not store above 25 °C.

Once opened, use within 6 months stored below 25 °C and 60 ± 5% RH.

Special warning

Special warnings for each target species: It is recommended to treat all animals in a pen. Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness. To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required. Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

As with any antiparasiticide, frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

Special precautions for use in animals: None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: People with known sensitivity to the active substance or any of the excipients should avoid contact with this product. Avoid skin and eye contact with the product. Wash any splashes from skin or eyes immediately with water. Do not eat, drink or smoke while using the product.

Other precautions: The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both persistent (half-life >1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle:

Veal calves: Do not use in veal calves.

Dairy calves: Do not use in dairy calves weighing more than 80 kg bodyweight.

In order to prevent any adverse effects on plants and possible contamination of groundwater manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Beef calves: Do not use in beef calves weighing more than 150 kg bodyweight.

Fattening calves: Do not use in fattening calves less than 3 months old.

Do not use in fattening calves weighing more than 150 kg bodyweight

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Pregnancy and lactation: Not applicable.

Interactions: None known. In pigs, interactions in combination with iron supplement do not exist.

Overdose: A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance. No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

Incompatibilities: In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Other information

Properties: Toltrazuril is a triazinon derivative. It acts against coccidia of the genus *Isospora* and *Eimeria*. It is active against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

Cattle: After oral administration toltrazuril is slowly absorbed. The maximal plasma concentration (C_{max} = 36.6 mg/l) was observed between 24 and 48 hours (geometric mean 33.9 hours) following oral administration. The elimination of toltrazuril is slow with a terminal half-life time of approximately 2.5 days (64.2 hours). The main metabolite is characterised as toltrazuril sulfone. The major route of excretion is via the faeces.

Pigs: After oral administration toltrazuril is slowly absorbed with a bioavailability of $\geq 70\%$. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with a half-life elimination time around 3 days. The major route of excretion is via the faeces.

Sheep: After oral administration toltrazuril is slowly absorbed in mammals. The main metabolite is characterised as toltrazuril sulfone. The maximal plasma concentration (C_{max} = 62 mg/L) was observed 2 days following oral administration. The elimination of toltrazuril is slow with an elimination half-life time of approximately 9 days. The major route of excretion is via the faeces.

Pack sizes: Bottles of 250 ml or 1 L. Not all pack sizes may be marketed.

For animal treatment only. To be prescribed and treated by registered veterinary surgeons only.

Name and address of the manufacturer:

Industrial Veterinaria, S.A.

C/ Esmeralda, 19

Esplugues de Llobregat, 08950

Barcelona, Spain

Marketing authorisation holder:

YENHER AGRO PRODUCTS SDN. BHD. (222455-P)

1628 Jalan IKS Simpang Ampat 1,

Taman IKS Simpang Ampat.

14100 Simpang Ampat, S.P.S.,

Penang, Malaysia.

Date of revision of package insert: February 2020