

Veterinary Package Insert

LSTYVET 50% W/W PREMIX POWDER (50% Tylosin Phosphate)

PRODUCT DESCRIPTION

White to pale yellow colour powder, which contains 500mg of tylosin phosphate per gram.

PHARMACODYNAMICS & PHARMACOKINETICS

Tylosin has the same mechanism of antibiotic action as other macrolides, that is, binding to the 50S ribosome fraction, resulting in inhibition of protein synthesis. Tylosin has a dominant bacteriostatic activity.

Tylosin is widely distributed in the body after absorption but does not cross the blood-brain barrier. It is eliminated by the urinary tract and bile unchanged. The half-life of elimination of tylosin is of the order of one to two hours depending on the animal species.

INDICATION

For tylosin-sensitive diseases in poultry and pigs.

Poultry:

Prevention in an infected environment and treatment of avian mycoplasmosis.

Chickens:

Prevention in infected medium of necrotic enteritis due to *Clostridium perfringens*.

Pigs:

Prevention in an infected environment of haemorrhagic enteritis.

Prevention in infected environment of enzootic pneumonia.

TARGET SPECIES

Poultry, chickens and pigs.

RECOMMENDED DOSAGE AND ADMINISTRATION

In poultry:

Prevention in infected medium and treatment of avian mycoplasmosis:

50 to 100 mg of tylosin (110.7 to 221.4mg of product) per kg of live weight and per day orally for 2 to 5 days.

For a daily dietary intake of 100 g of feed per kg of live weight, this dosage corresponds to 500 to 1000 ppm of tylosin in the feed, ie an incorporation rate of 11.07 kg to 22.14 kg of medicated premix per ton of feed.

In chickens:

Prevention in infected medium of necrotic enteritis due to *Clostridium perfringens*:

10 to 20 mg of tylosin (22.1 to 44.3mg of product) per kg of live weight and per day orally for 7 days.

For a daily dietary intake of 100 g of feed per kg of live weight, this dosage corresponds to 100 ppm to 200 ppm of tylosin in the feed, ie an incorporation rate of 2.21kg to 4.43kg of medicated premix per ton of food.

In pigs:

Prevention in an infected environment of haemorrhagic enteritis and prevention in an infected environment of enzootic pneumonia:

5 mg of tylosin (11.1mg of product) per kg of live weight and per day orally for 3 weeks.

For a daily dietary intake of 50 g of feed per kg of live weight, this dosage corresponds to 100 ppm of tylosin in the feed, ie an incorporation rate of 2.21 kg of medicated premix per ton of feed.

CONTRAINDICATIONS

Do not use in animals known to be sensitive to tylosin or other macrolides.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals

This medicated premix is intended for the manufacture of medicated feeds and can not be used as is; the rate of incorporation of the premix into the feed cannot be less than 2.21kg / tonne.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Macrolides may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to tylosin and other macrolides can be crossed. Allergic reactions to these substances can be serious.

In case of hypersensitivity, avoid contact with the product.

If skin erythema develops, take medical advice.

If edema of the face, lips, eyes, or if breathing is difficult, seek immediate medical attention.

Avoid inhalation of dust and avoid contact with skin and eyes when handling the product.

In case of accidental projection into the eyes, rinse thoroughly with clear water.

Wash hands after use of the product.

INTERACTIONS WITH OTHER MEDICATIONS

Do not administer at the same time as bactericidal antibiotics.

PREGNANCY AND LACTATION

Studies in laboratory animals have not shown any teratogenic or embryotoxic effects or consequences on the fertility of animals.

The safety of the specialty has not been studied in pregnant pigs. The use of the specialty in the pregnant pig should be the subject of a benefit / risk assessment by the veterinarian.

ADVERSE EFFECTS

It has been reported in pigs the possibility of digestive side effects, with edema of the rectal mucosa, anal prolapse with pruritus, erythema and diarrhea.

OVERDOSE & TREATMENT

Not known.

WITHDRAWAL PERIOD

Meat and offal:

Pigs: 0 days.

Chickens: 0 days.

Poultry other than chickens: 2 days.

Eggs:

Chickens: 0 days.

Poultry other than hens: in the absence of a waiting time for eggs do not use in laying birds producing egg for consumption (4 weeks before the start of laying and during it).

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in food µg/kg
Tylosin	Muscle, liver, kidney, skin + fat (Porcine)	100
	Muscle, liver, kidney, skin + fat (Poultry)	100

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

1000gm

LIM SENG PHARMACHEM SDN. BHD.

(product registration holder)

2692, Tingkat Perusahaan Enam,
Kawasan Perusahaan Prai Mk 1,
13600 Prai, Pulau Pinang.

THYE PHARMA SDN. BHD.

(manufacturer)

No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised January 2020