

Lyzodase Oral Liquid

Lysozyme Hydrochloride 20 mg /5 ml, Pyridoxine Hydrochloride (Vit. B6) 10mg/5ml

Methylparaben 0.05%w/v, propylparaben 0.005%w/v

No added sugar oral liquid.

Product Description: A clear, thick, viscous and sweet liquid with banana flavour.

Pharmacodynamics:

Lysozyme is an enzyme having a mucopolysaccharide lysing action. It enhances the effect of antibiotics to destroy or weaken cell bacteria. Lysozyme also accelerates immuno-bacteriolysis and activates the phagocytosis activity of leukocytes to destroy bacteria and viruses. Lysozyme is widely distributed in tears, nasal discharge, saliva and also breast milk. Lysozyme extracted from egg-white is used clinically for the intensification of natural defence mechanisms, restoration of damaged tissue and anti-inflammation.

The body needs pyridoxine hydrochloride in order to make cells of the immune system. Pyridoxine hydrochloride helps to improve immune response to the increase in production of antibodies and also helps in communicative interactions between cytokines and chemokines.

Pharmacokinetics:

In healthy men receiving a single 900 mg (potency) oral dose of lysozyme chloride, the peak plasma concentration occurred about 30 minutes to 1 hour after administration. Animal studies - Absorption and Distribution: following administration of [1]-labelled egg white lysozyme (2mg/kg) into the intestinal tract of rats, peak serum and lymphatic concentration occurred at 30 minutes.

Pyridoxine hydrochloride is absorbed rapidly from the upper intestine regardless of the size of the dose given. Absorption may also occur from the ileum and to a small extent from the colon. There is a linear relationship between oral dose and the amount absorbed in normal animals and in those in which the distal small intestine has been resected. These observations suggest the possibility that pyridoxine may be absorbed by diffusion. Pyridoxine is rapidly converted in the liver to pyridoxine phosphate, pyridoxal phosphate and pyridoxamine phosphate via oxidation, this causes the release of pyridoxal and so pyridoxal phosphate to the general circulation where it reaches other organs chiefly as circulating pyridoxal. In urine, pyridoxine was excreted primarily unchanged with a small amount of only one metabolite, most likely 4-pyridoxic acid.

Indication: A soothing preparation for relief of sore throat.

Recommended dose:

Children 6 – 12 years old: 5 ml

Adults: 10 ml

Sip the dose slowly. Repeat dose as required up to 4 times daily.

If the symptoms worsen or do not improve within 7 days, consult your doctor.

Contraindications:

Patients hypersensitive to egg-white or vitamin B6.

Patients with a history of hypersensitivity to any ingredients of the product.

Warnings and Precautions:

Patients with an allergic predisposition manifested as atopic dermatitis, bronchial asthma, drug allergy or food allergy. Abuse of pyridoxine may be responsible for sensory neuropathy that occurs with high doses of pyridoxine (> 200 mg/day) for a long period of time (months to years). This is usually reversible when the treatment is stopped.

Pregnancy and Lactation: It is best not to use this medicine during pregnancy and breastfeeding.

Lyzodase Oral Liquid

Lysozyme Hydrochloride 20 mg /5 ml, Pyridoxine Hydrochloride (Vit. B6) 10mg/5ml

Methylparaben 0.05%w/v, propylparaben 0.005%w/v

No added sugar oral liquid.

Side effects:

May cause diarrhoea, anorexia, stomach discomfort, nausea and vomiting. May cause hypersensitivity reactions such as skin rashes. In rare case, shock may occur with administration of this drug. Therefore, patients should be carefully monitored.

Symptoms and Treatment of Overdose:

Symptoms: diarrhoea, anorexia, gastric discomfort.

Treatment: Stop administration.

Storage: Store below 30°C.

Shelf-life: 3 years

Packing: 90ml amber PET bottle with HDPE cap

Product Registration Holder:

Product registration holder: Vismedic Sdn Bhd (890098-W)

28B, Jalan BP 6/6, Bandar Bukit Puchong, 47120 Puchong, Selangor.

Manufacturer:

AV Manufacturing Sdn. Bhd (667760-V)

Lot 10621 (PT16700), Jalan Permata 2, Arab-Malaysian Industrial Park,
71800 Nilai, Negeri Sembilan, Malaysia.

Dec 2023

Ver. 02