

VENAFIX

Diosmin and Hesperidin Tablets

COMPOSITION:

Each film coated tablet contains:

Diosmin BP 450mg
Hesperidin 50mg

Product Description:

Light brown coloured, oblong shaped, biconvex, film-coated tablets plain on both sides.

Pharmacodynamics/ Pharmacokinetics:

Pharmacotherapeutic group: Venotonic and vascular protector.

ATC code: C05CA53.

Pharmacology

It is active upon the return vascular system in the following way:

- it reduces venous distensibility and stasis,
- in the microcirculation, it normalises capillary permeability and increases capillary resistance.

Pharmacokinetics

Absorption

Following oral administration, diosmin is rapidly hydrolyzed in the intestine by intestinal flora and absorbed as its aglycone derivative, diosmetin. Oral bioavailability is approximately 57.9%.

Distribution

Diosmetin has a volume of distribution of 62.1 L indicating a

broad distribution into tissues.

Biotransformation

Diosmetin is extensively metabolized to phenolic acids or to its glucuronide derivatives of aglycone that are eliminated in urine.

The major metabolite found in urine is m-hydroxyphenylpropionic acid which is mainly eliminated in its conjugated form. Metabolites found in small quantities include phenolic acids corresponding to 3-hydroxy-4 methoxybenzoic acid, 3-metoxy-4-hydroxyphenylacetic acid and 3,4-dihydroxybenzoic acid.

Elimination

The elimination half-life of diosmetin showed a mean value of 31.5 hours, ranging between 26 and 43 hours.

Indication

- Treatment of symptoms related to venolymphatic insufficiency (heavy legs, pain, early morning restless legs).
- Treatment of functional symptoms related to acute hemorrhoidal attack.


Recommended Dosage:

- Usual dosage: 2 tablets daily in two divided doses, midday and evening at meal times.
- Acute hemorrhoidal attack: 6 tablets per day for the first 4 days, then 4 tablets per day for 3 days.

Route of Administration: Oral use

Contraindications:

Hypersensitivity to the active substance or to any of the excipients.



Warnings and Precautions:

Jauhkan daripada capaian kanak-kanak.

Keep out of reach of children.

Please consult your pharmacist/doctor before taking this product.

The administration of this product for the symptomatic treatment of acute hemorrhoids does not preclude treatment for other anal conditions. If symptoms do not subside promptly, a proctological examination should be performed and the treatment should be reviewed.

Interactions with Other Medicaments:

No interaction studies have been performed. However and taking into consideration the huge post marketing experience on the product, no drug interaction has been reported to date.

Statement on usage during pregnancy and lactation:

Pregnancy:

No teratogenic effects have been shown in several studies and no adverse effect have been reported in human. As a precautionary measure, it is preferable to avoid the use of product during pregnancy.

Breast-feeding:

In the absence of data on excretion in milk, treatment should be avoided during breast-feeding.

Adverse Effects/ Undesirable Effects:

Tabulated list of adverse reactions

The following adverse effects or events have been reported and are ranked using the following frequency: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000); not known (cannot be estimated form the available data).

System Organ Class	Frequency	Preferred Term
Nervous system disorders	Rare	Dizziness
		Headache
		Malaise

Gastrointestinal disorders	Common	Diarrhoea
		Dyspepsia
		Nausea, Vomiting
	Uncommon	Colitis
Skin and subcutaneous tissue disorders	Not known*	Abdominal pain
	Rare	Pruritus
		Rash
		Urticaria
	Not known*	Isolated face, lip, eyelid oedema. Exceptionally Quincke's oedema

* Post-marketing experience

Overdose and Treatment:

No case of overdose with Diosmin and Hesperidin tablet has been reported.

Storage:

Store at temperatures not exceeding 30°C.
Keep all medicines out of children's reach.

AVAILABILITY:

Alu / PVC blister pack of 3x10's & 10x10's

Date of revision of PI
14/11/2024

XL

Manufactured by :
XL LABORATORIES PVT. LTD.
E-1223, Phase-I Extn., (Ghatal) RIICO Indl. Area, Bhiwadi-301019, Rajasthan, India

Product Registration Holder:
Healol Pharmaceuticals Sdn.Bhd.
74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju,
53300 Kuala Lumpur, Malaysia.

