



What is ORACUE?

Description, Dosage form and Packaging

ORACUE is a translucent colourless gel containing Choline Salicylate 8.71%w/w and Cetalkonium Chloride 0.01%w/w. Contains Ethanol 33.46% w/w. This product is available in pack size of 10g tube.

What is ORACUE used for?

ORACUE helps to relieve pain, inflammation, and to treat minor infection due to common mouth ulcers, cold sores, denture and sore spots, as well as mouth ulcers and sore spots due to orthodontic devices. To help to fight minor mouth infection and healing of sore spots and ulcers due to dentures and orthodontic devices. Oracue helps to relieve and soothe teething.

How ORACUE works?

ORACUE contains non-steroidal anti-inflammatory (NSAID) and anti-infective substances.

Choline Salicylate is a NSAID that reduces pain and inflammation. Cetalkonium Chloride is an anti-infective agent that fights against microbial growth.

Pharmacodynamics

Choline salicylate is the choline salt of salicylic acid and its pharmacology is essentially that of salicylic acid. It has exhibited anti-inflammatory, analgesic and antipyretic actions in animal models, and is taken orally or is applied topically in man for the relief of pain and inflammation. Like salicylic acid it has no antithrombotic activity and shows a low potential for production of gastrointestinal injury when given by the oral route. The pharmacological actions of choline salicylate are thought to be primarily mediated through inhibition of prostaglandin production, although effects on leukotriene pathways, kinin release and nerve conduction have been proposed. Cetalkonium chloride is a quaternary ammonium antimicrobial agent, being bactericidal towards both Gram positive and Gram negative organisms, but with preferences for the former.

Pharmacokinetics

Choline salicylate is absorbed from the gut and is likely to be absorbed across mucous membrane such as all buccal mucosa. Metabolism of salicylic acid is by glycine and phenolic or acyl glucuronate conjugation with small amounts undergoing hydroxylation. The plasma half-life of salicylic acid is 2-4 hours. Both metabolites and a small amount of intact salicylic acid are excreted, mainly in the urine. Salicylic acid is highly protein bound (80-90%) and although it has a low apparent volume of distribution of around 0.15 l/kg, it is widely distributed throughout most tissues and extracellular water.

How to use ORACUE?

ORACUE gel is for topical oral mucosa use only. Always wash your hand before applying the gel. Make sure the hand is clean.

Babies (4 month-old and above) and Children

With clean finger, massage approximately one half centimeter of ORACUE gel onto the sore area. This can be repeated after 3 hours.

Not to exceed 6 doses in 24 hours. Not to be used in babies less than 4 month-old.

Adults

Massage approximately one centimeter of ORACUE gel onto the sore area. This can be repeated after 3 hours.

Denture Sores

Massage approximately one centimeter of ORACUE gel onto the sore area and leave at least 30 minutes before reinserting the dentures.

Do not apply directly to the dentures.

OR0100-001

Precaution

Please always consult your doctor, dentist or pharmacist before using ORACUE gel.

Tell your doctor, dentist or pharmacist if you:

- are allergic to chlorine salicylates, cetalkonium chloride, ethanol or glycerin
- taking medicines for blood clotting and/or gout treatment
- have stomach or intestinal bleeding
- have blood clotting disorders
- are pregnant or breast-feeding

Do not exceed the stated dose. If symptoms persist for more than seven days, consult your doctor or dentist. Keep out of the reach of children. Contains ethanol.

Pregnancy and Lactation

Pregnancy

There is clinical evidence of the safety of salicylates in pregnancy, but may prolong bleeding and lead to maternal and neonatal bleeding, and are best avoided at term.

Lactation

Salicylates are excreted at low concentrations in breast milk, but are unlikely to adversely affect the infant.

Interactions With Other Medicaments

Choline Salicylate may enhance the effect of anticoagulants and inhibit the action of uricosurics.

Contraindication

Not to be used in babies less than 4 month-old.

Contraindication for patients suffering from active peptic ulceration or known to be allergic to salicylates.

Does ORACUE have side effects?

Asthma attacks may occasionally occur in people who are sensitive to salicylates. Salicylates may precipitate bronchospasm and induce asthma attacks in susceptible patients.

As this product containing glycerin, some people may experience headache, stomach upset and diarrhea.

If you experience any symptoms persistently for more than 7 days, please consult your doctor, dentist or pharmacist.

What to do if you have used ORACUE more than the recommended dose?

In case of overdose, seek medical attention immediately.

Overdose may lead to salicylate toxicity. Symptoms of salicylate poisoning are such as nausea, vomiting, tinnitus and excessive sweating. The usual procedure for salicylate overdosage should be followed, including general supportive measures and gastric lavage if necessary.

How should you keep ORACUE gel?

Store below 30 °C. Keep out of reach of children.

The expiry date of this product is indicated on the packaging.

Revision date: 15-May-2020.

Manufacturer and Product Registration Holder (in Malaysia)
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

