

Pigfen 40 mg/g premix for medicated feeding stuff for pigs

Keep out of reach of children.

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Per gram:
Fenbendazole 40 mg

PRODUCT DESCRIPTION

Pigfen 40 mg/g premix for medicated feeding stuff for pigs represents off-white to light yellow granules.

PHARMACODYNAMICS

Fenbendazole is an anthelmintic belonging to the benzimidazole-carbamate group. It acts by binding to beta-tubulin, thereby inhibiting the polymerisation of tubulin to microtubules and subsequently interfering with energy metabolism. The anthelmintic affects both adult and immature stages of *Ascaris suum*.

PHARMACOKINETICS

Fenbendazole is only partly absorbed after oral administration and is then metabolised in the liver. Body clearance of fenbendazole in serum after intravenous administration to pigs at a dose rate of 1 mg/kg was 1.36 L/h/kg, volume of distribution at steady state was 3.35 L/kg and the mean residence time was 2.63 hours. After oral administration at a dose rate of 5 mg/kg the peak plasma concentration of fenbendazole was 0.07 µg/ml, the T_{max} was 3.75 hours and the mean residence time was 15.15 hours. The bioavailability was 27.1 %. Oxfendazole was the major plasma metabolite i.e. 2/3 of the total AUC.

Fenbendazole and its metabolites are distributed throughout the body and high concentrations can be found in the liver.

The elimination of fenbendazole and its metabolites occurs primarily via the faeces (>90%) and to a small extent in the urine and milk.

Fenbendazole is metabolised to its sulfoxide, then to sulphone and amines.

INDICATION(S)

Treatment of pigs infected with *Ascaris suum* (adult, intestinal and migrating larval stages) susceptible to fenbendazole.

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

The product is suitable for herd medication of pigs. Administer at a dose rate of 5 mg fenbendazole per kg bodyweight.

May be administered to pigs either as a single dose of 5 mg/kg (migrating larval, intestinal larval and adult stages) or by divided dose of 0.72 mg/kg over 7 days (intestinal larval and adult stages) or 0.36 mg/kg over 14 days (intestinal larval and adult stages).

Single dose treatment

Use the following formula to calculate how much product to add per tonne of feed:

$$\text{Kg} = \frac{\text{Bodyweight (kg)}}{\text{Powder per tonne (Daily feed intake (kg) x 8)}}$$

7 day treatment

The standard dose rate can be divided and administered in feed over 7 days. Use the following formula to calculate how much product to add per tonne of feed:

$$\text{Kg} = \frac{\text{Bodyweight (kg)}}{\text{Powder per tonne (Daily feed intake (kg) x 56)}}$$

14 day treatment

The standard dose rate can be divided and administered in feed over 14 days. Use the following formula to calculate how much product to add per tonne of feed:

$$\text{Kg} = \frac{\text{Bodyweight (kg)}}{\text{Powder per tonne (Daily feed intake (kg) x 112)}}$$

MODE OF ADMINISTRATION

For oral administration. In-feed use.

To avoid under dosing, all incorporation rate calculations as presented above should be based on the heaviest pig in the group. For incorporation into dry feed at the registered mill.

A manufacturer who is approved to incorporate veterinary medicinal products, or premixtures containing such products, directly at any concentration, must be responsible for mixing when incorporation is less than 2 kg per tonne for final feed. To ensure adequate distribution of the product in the final feed it is recommended to premix the product at a ratio of 1:10 with feed ingredients before blending into the final feed. If the premix is used for supplementation of pelleted feed, the pelleting temperature should not exceed 85 °C. Not to be mixed in liquid feed.

CONTRAINDICATIONS

Do not use in known cases of hypersensitivity to the active substance, other benzimidazoles or any of the excipients.

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.

Under dosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any). Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used. Inappetent animals should be treated individually.

Special precautions for use in animals:

None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This product may cause eye irritation. Avoid contact with skin and/or eyes. Do not eat or smoke during handling the premix or the medicated feed. When handling or mixing, care should be taken to avoid direct contact with the skin and eyes, and inhalation of dust, by wearing goggles, impervious gloves and a disposable half-mask respirator conforming to European Standard EN149 or a non- disposable respirator to European Standard EN 140 with a filter to EN 143. Wash hands after use. In case of skin and/or eye contact, immediately rinse with plenty of water. **OTHER PRECAUTIONS:** The veterinary medicinal product should not be allowed to enter surface waters as it has harmful effects on aquatic organisms.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

Exacerbation of paracetamol hepatotoxicity by fenbendazole cannot be excluded

PREGNANCY AND LACTATION

The product can be safely administered to pregnant animals.

The safety of the veterinary medicinal product has not been established during lactation. Use only according to the benefit-risk assessment by the responsible veterinarian.

ADVERSE REACTIONS

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

OVERDOSE

Pigfen administered as a single 25 mg fenbendazole/kg dose for three consecutive days did not produce any clinically apparent adverse reactions in pigs. In addition, it has been shown that administration of non-formulated fenbendazole at a dose of 2000 mg/kg for 14 consecutive days was well tolerated in pigs.

INCOMPATIBILITIES

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products

WITHDRAWAL PERIOD

Meat and offal: 4 days

DISPOSAL

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. The product should not enter water courses as this may be dangerous for aquatic organisms.

STORAGE CONDITIONS

Store in the original container to protect from light. Store in a dry place. Store below 30°C.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month. Shelf life after first opening the container: 3 months. Shelf life after incorporation into meal or pelleted feed: 3 months when stored below 25°C and 4 weeks when stored below 40°C.

Medicated feed (mashed and pelleted): no special storage precautions.

PACKAGING AVAILABLE

The product is distributed in multiple-layer paper bag with internal aluminium/polyethylene layer and with sutured crimp of 20 kg, as well as in polyethylene/aluminium foil/polyethylene terephthalate zipper bag of 1, 2 and 5 kg.

Not all pack sizes may be marketed.

MANUFACTURER

BIOVET Joint Stock Company
39 Petar Rakov Street, 4550 Peshtera, Bulgaria

MARKETING AUTHORISATION HOLDER

Pahang Pharmacy Sdn. Bhd. (10826-K)
Lot 5979 Jalal Teratai, 5 ½ Mile Off Jalan Meru, 41050 Klang, Selangor Darul Ehsan, Malaysia

DATE OF REVISION

December, 2018.

