

Prescribing Information

For the use of Registered Medical Practitioner,
Hospital or a Laboratory Only.

ROCUTROY

Rocuronium Bromide 10mg/ml
Solution for Injection or Infusion

COMPOSITION :

Each ml contains :
Rocuronium Bromide EP 10 mg
Water for Injections BP q.s.
Other excipients, see EXCIPIENTS

DESCRIPTION :

ROCUTROY Rocuronium Bromide 10mg/ml Solution for Injection or Infusion is a clear, colourless to slightly yellow coloured solution filled in a 5ml Flint tubular clear glass vial USP Type-I with 20 mm grey coloured bromo butyl rubber stopper and 20mm light pink flip off seal packed in printed carton with printed insert.

After dilution: Clear colourless to yellowish colour solution

INDICATIONS :

Rocutroy is indicated as an adjunct to general anaesthesia to facilitate endotracheal intubation, to provide skeletal muscle relaxation and to facilitate mechanical ventilation in adults, children and infants from one month of age. Rocutroy is also indicated as an adjunct in the intensive care unit (ICU) to facilitate mechanical ventilation as part of Rapid Sequence Induction, however, this has not been studied in infants and children.

DOSAGE AND ADMINISTRATION :

Dosage

Like other neuromuscular blocking agents, Rocutroy should only be administered by, or under supervision of, experienced clinicians who are familiar with the action and use of these drugs.

As with other neuromuscular blocking agents, the dosage of Rocutroy should be individualized in each patient. The method of anaesthesia and the expected duration of surgery, the method of sedation and the expected duration of mechanical ventilation, the possible interaction with other drugs that are administered concomitantly, and the condition of the patient should be taken into account when determining the dose.

The use of an appropriate neuromuscular monitoring technique is recommended for the evaluation of neuromuscular block and recovery.

Inhalational anaesthetics do potentiate the neuromuscular blocking effects of Rocutroy. This potentiation however, becomes clinically relevant in the course of anaesthesia, when the volatile agents have reached the tissue concentrations required for this interaction. Consequently, adjustments with Rocutroy should be made by administering smaller maintenance doses at less frequent intervals or by using lower infusion rates of Rocutroy during long lasting procedures (longer than 1 hour) under inhalational anaesthesia (see Interaction with other medicinal products section and other forms of interaction).

Risk of Medication Errors: Accidental administration of neuromuscular blocking agents may result in serious adverse events, including fatal outcomes. Store ROCUTROY with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product [see Warnings and precautions for use] In adult patients the following dosage recommendations may serve as a general guideline for tracheal intubation and muscle relaxation for short to long lasting surgical procedures and for use in the intensive care unit.

Surgical Procedures

Tracheal intubation : The standard intubating dose during routine anaesthesia is 0.6 mg.kg⁻¹ rocuronium bromide, after which adequate intubation conditions are established within 60 seconds in nearly all patients. A dose of 1.0 mg.kg⁻¹ of rocuronium bromide is recommended for facilitating tracheal intubation conditions during rapid sequence induction of anaesthesia, after which adequate intubation conditions are established within 60 seconds in nearly all patients. If a dose of 0.6 mg.kg⁻¹ rocuronium bromide is used for rapid sequence induction of anaesthesia, it is recommended to intubate the patient 90 seconds after administration of rocuronium bromide.

For use of rocuronium bromide during rapid sequence induction of anaesthesia in patients undergoing Cesarean section reference is made to section Pregnancy and lactation.

Maintenance dosing : The recommended maintenance dose is 0.15 mg.kg⁻¹ rocuronium bromide; in the case of long-term inhalational anaesthesia, this should be reduced to 0.075-0.1 mg.kg⁻¹ rocuronium bromide. The maintenance doses should best be given when twitch height has recovered to 25% of control twitch height, or when 2 to 3 responses to train of four stimulation are present.

Continuous infusion : If rocuronium bromide is administered by continuous infusion, it is recommended to give a loading dose of 0.6 mg.kg⁻¹ rocuronium bromide and, when neuromuscular block starts to recover, to start administration by infusion. The infusion rate should be adjusted to maintain twitch response at 10% of control twitch height or to maintain 1 to 2 responses to train of four stimulation. In adults under intravenous anaesthesia, the infusion rate required to maintain neuromuscular block at this level ranges from 0.3-0.6 mg.kg⁻¹.h⁻¹ and under inhalational anaesthesia the infusion rate ranges from 0.3-0.4 mg.kg⁻¹.h⁻¹. Continuous monitoring of neuromuscular block is recommended since infusion rate requirements vary from patient to patient and with the anaesthetic method used.

Pediatric patients : For infants (28 days-23 months), children (2-11 years) and adolescents (12-18 years) the recommended intubation dose during routine anaesthesia and maintenance dose are similar to those in adults.

For continuous infusion in paediatrics, the infusion rates, with exception of children, are the same as for adults. For children higher infusion rates might be necessary. For children the same initial infusion rates as for adults are recommended and this should be adjusted to maintain twitch response at 10% of control twitch height or to maintain 1 or 2 responses to train of four stimulation during the procedure.

There are insufficient data to support dose recommendations for the use of rocuronium bromide in neonates (0-1 month).

The experience with rocuronium bromide in rapid sequence induction in Pediatric patients is limited. Rocuronium bromide is therefore not recommended for facilitating tracheal intubation conditions during rapid sequence induction in Pediatric patients.

Geriatric patients and patients with hepatic and/or biliary tract disease and/or renal failure : The standard intubation dose for geriatric patients and patients with hepatic and/or biliary tract disease and/or renal failure during routine anaesthesia is 0.6 mg.kg⁻¹ rocuronium bromide. A dose of 0.6 mg.kg⁻¹ should be considered for rapid sequence induction of anaesthesia in patients in which a prolonged duration of action is expected. Regardless of the anaesthetic technique used, the recommended maintenance dose for these patients is 0.075-0.1 mg.kg⁻¹ rocuronium bromide, and the recommended infusion rate is 0.3-0.4 mg.kg⁻¹.h⁻¹ (see Continuous infusion) (See also section warnings and precautions for use).

Overweight and obese patients : When used in overweight or obese patients (defined as patients with a body weight of 30% or more above ideal body weight) doses should be reduced taking into account ideal body weight.

Intensive Care Procedures

Tracheal intubation : For tracheal intubation, the same doses should be used as described above under surgical procedures.

Maintenance dosing : The use of an initial loading dose of 0.6 mg.kg⁻¹ rocuronium bromide is recommended, followed by a continuous infusion as soon as twitch height recovers to 10% or upon reappearance of 1 to 2 twitches to train of four stimulation. Dosage should always be titrated to effect in the individual patient. The recommended initial infusion rate for the maintenance of a neuromuscular block of 80-90% (1 to 2 twitches to TOF stimulation) in adult patients is 0.3-0.6 mg.kg⁻¹.h⁻¹ during the first hour of administration, which will need to be decreased during the following 6-12 hours, according to the individual response. Thereafter, individual dose requirements remain relatively constant.

A large between patient variability in hourly infusion rates has been found in controlled clinical studies, with mean hourly infusion rates ranging from 0.2-0.5 mg.kg⁻¹.h⁻¹ depending on nature and extent of organ failure(s), concomitant medication and individual patient characteristics. To provide optimal individual patient control, monitoring of neuromuscular transmission is strongly recommended. Administration up to 7 days has been investigated.

Special populations : Rocutroy is not recommended for the facilitation of mechanical ventilation in the intensive care in Pediatric and geriatric patients due to a lack of data on safety and efficacy.

Administration

Rocutroy is administered intravenously either as a bolus injection or as a continuous infusion (see section PRECAUTIONS FOR USE AND OTHER HANDLING).

CONTRAINDICATIONS :

Hypersensitivity to rocuronium or to the bromide ion or to any of the excipients.

WARNINGS AND PRECAUTIONS :

Since Rocuronium causes paralysis of the respiratory muscles, ventilatory support is mandatory for patients treated with this drug until adequate spontaneous respiration is restored. As with all neuromuscular blocking agents, it is important to anticipate intubation difficulties, particularly when used as part of a rapid sequence induction technique.

As with other neuromuscular blocking agents, residual neuromuscular blockade has been reported for rocuronium. In order to prevent complications resulting from residual neuromuscular blockade, it is recommended to extubate only after the patient has recovered sufficiently from neuromuscular block. Geriatric patients (65 years or older) may be at increased risk for residual neuromuscular block. Other factors which could cause residual neuromuscular blockade after extubation in the post-operative phase (such as drug interactions or patient condition) should also be considered. If not used as part of standard clinical practice, the use of a reversal agent (such as sugammadex or acetylcholinesterase inhibitors) should be considered, especially in those cases where residual neuromuscular blockade is more likely to occur.

High rates of cross-sensitivity between neuromuscular blocking agents have been reported. Therefore, where possible, before administering rocuronium, hypersensitivity to other neuromuscular blocking agents should be excluded. Rocuronium should only be used when absolutely essential in susceptible patients. Patients who experience a hypersensitivity reaction under general anaesthesia should be tested subsequently for hypersensitivity to other neuromuscular blockers.

Rocuronium may increase the heart rate.

In general, following long term use of neuromuscular blocking agents in the ICU, prolonged paralysis and/or skeletal muscle weakness has been noted. In order to help preclude possible prolongation of neuromuscular block and/or overdose it is strongly recommended that neuromuscular transmission is monitored throughout the use of neuromuscular blocking agents. In addition, patients should receive adequate analgesia and sedation. Furthermore, neuromuscular blocking agents should be titrated to effect in the individual patients by or under supervision of experienced clinicians who are familiar with their actions and with appropriate neuromuscular monitoring techniques. Myopathy after long term administration of other non-depolarising neuromuscular blocking agents in the ICU in combination with corticosteroid therapy has been reported regularly. Therefore, for patients receiving both neuromuscular blocking agents and corticosteroids, the period of use of the neuromuscular blocking agent should be limited as much as possible.

If suxamethonium is used for intubation, the administration of rocuronium should be delayed until the patient has clinically recovered from the neuromuscular block induced by suxamethonium.

Because rocuronium bromide is always used with other drugs and because of the risk of malignant hyperthermia during anaesthesia, even in the absence of known triggering factors, physicians should be aware of the early symptoms, confirmatory diagnosis and treatment of malignant hyperthermia prior to the start of anaesthesia. Animal studies have shown that rocuronium bromide is not a triggering factor for malignant hyperthermia. Rare cases of malignant hyperthermia with rocuronium have been observed thru post-marketing surveillance; however, the causal association has not been proven.

The following conditions may influence the pharmacokinetics and/or pharmacodynamics of Rocuronium :

Hepatic and/or biliary tract disease and renal failure : Because rocuronium is excreted in urine and bile, it should be used with caution in patients with clinically significant hepatic and/or biliary diseases and/or renal failure. In these patient groups prolongation of action has been observed with doses of 0.6 mg/kg rocuronium bromide.

Prolonged circulation time : Conditions associated with prolonged circulation time such as cardiovascular disease, old age and oedematous state resulting in an increased volume of distribution, may contribute to a slower onset of action. The duration of action may also be prolonged due to a reduced plasma clearance.

Neuromuscular disease : Like other neuromuscular blocking agents, Rocuronium should be used with extreme caution in patients with a neuromuscular disease or after poliomyelitis since the response to neuromuscular blocking agents may be considerably altered in these cases. The magnitude and direction of this alteration may vary widely. In patients with myasthenia gravis or with the myasthenic (Eaton-Lambert) syndrome, small

doses of rocuronium may have profound effects and rocuronium should be titrated to the response.

Hypothermia : In surgery under hypothermic conditions, the neuromuscular blocking effect of rocuronium is increased and the duration prolonged.

Obesity : Like other neuromuscular blocking agents, rocuronium may exhibit a prolonged duration and a prolonged spontaneous recovery in obese patients when the administered doses are calculated on actual body weight.

Burns : Patients with burns are known to develop resistance to non-depolarising neuromuscular blocking agents. It is recommended that the dose is titrated to response.

Conditions which may increase the effects of Rocuronium : Hypokalaemia (e.g. after severe vomiting, diarrhoea and diuretic therapy), hypermagnesaemia, hypocalcaemia (after massive transfusions), hypoproteinaemia, dehydration, acidosis, hypercapnia, cachexia.

Severe electrolyte disturbances, altered blood pH or dehydration should therefore be corrected when possible.

INTERACTION WITH OTHER MEDICAMENTS :

The following drugs have been shown to influence the magnitude and/or duration of action of non-depolarising neuromuscular blocking agents.

Effect of other drugs on Rocuronium

Increased effect:

- Halogenated volatile anaesthetics potentiate the neuromuscular block of rocuronium. The effect only becomes apparent with maintenance dosing. Reversal of the block with acetylcholinesterase inhibitors could also be inhibited.
- After intubation with suxamethonium.
- Long-term concomitant use of corticosteroids and rocuronium in the ICU may result in prolonged duration of neuromuscular block or myopathy.

Other drugs:

- antibiotics: aminoglycoside, lincosamide and polypeptide antibiotics, acylamino-penicillin antibiotics.
- diuretics, quinidine and its isomer quinine, magnesium salts, calcium channel blocking agents, lithium salts, local anaesthetics (lidocaine i.v, bupivacaine epidural) and acute administration of phenytoin or β -blockading agents.

Recurarisation has been reported after post-operative administration of: aminoglycoside, lincosamide, polypeptide and acylamino-penicillin antibiotics, quinidine, quinidine and magnesium salts.

Decreased effect :

- Prior chronic administration of phenytoin or carbamazepine.
- Calcium chloride, potassium chloride.
- Protease inhibitors (gabexate, ulinastatin).

Variable effect:

- Administration of other non-depolarising neuromuscular blocking agents in combination with rocuronium may produce attenuation or potentiation of the neuromuscular block, depending on the order of administration and the neuromuscular blocking agent used.
- Suxamethonium given after the administration of rocuronium may produce potentiation or attenuation of the neuromuscular blocking effect of Rocutroy.

Effect of Rocuronium on other drugs

Rocuronium combined with lidocaine may result in a quicker onset of action of lidocaine.

Paediatric population : No formal interaction studies have been performed. The above mentioned interactions for adults and their special warnings and precautions for use should be taken into account for paediatric patients.

PREGNANCY AND LACTATION :

Pregnancy : For rocuronium bromide, no clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, partition or postnatal development. Caution should be exercised when prescribing rocuronium to pregnant women.

Caesarean section : In patients undergoing Caesarean section, rocuronium can be used as part of a rapid sequence induction technique, provided no intubation difficulties are anticipated and a sufficient dose of anaesthetic agent is administered or following suxamethonium facilitated intubation. However, rocuronium, administered in doses of 0.6 mg/kg may not produce adequate conditions for intubation until 90 seconds after administration. This dose has been shown to be safe in parturients undergoing Caesarean section. Rocuronium does not affect Apgar score, foetal muscle tone or

cardiorespiratory adaptation. From umbilical cord blood sampling it is apparent that only limited placental transfer of rocuronium bromide occurs which does not lead to the observation of clinical adverse effects in the newborn.

Note 1: doses of 1.0 mg/kg have been investigated during rapid sequence induction of anaesthesia, but not in Caesarean section patients. Therefore, only a dose of 0.6 mg/kg is recommended in this patient group.

Note 2: Reversal of neuromuscular block induced by neuromuscular blocking agents may be inhibited or unsatisfactory in patients receiving magnesium salts for toxemia of pregnancy because magnesium salts enhance neuromuscular blockade. Therefore, in these patients the dosage of rocuronium should be reduced and be titrated to twitch response.

Breast-feeding : It is unknown whether rocuronium bromide is excreted in human breast milk. Animal studies have shown insignificant levels of rocuronium bromide in breast milk.

Insignificant levels of rocuronium bromide were found in the milk of lactating rats. There are no human data on the use of Rocutroy during lactation. Rocutroy should be given to lactating women only when the attending physician decides that the benefits outweigh the risks. After the administration of a single dose, it is recommended to abstain from next breastfeeding for five elimination half-lives of rocuronium, i.e. for about 6 hours.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES :

Since rocuronium is used as an adjunct to general anaesthesia, the usual precautionary measures after a general anaesthesia should be taken for ambulatory patients.

ADVERSE EFFECTS :

The most common undesirable effects are pain/reaction around injection site, changes in vital functions and prolonged neuromuscular block.

Immune system disorders

Very rare: Hypersensitivity, Anaphylactic reaction, Anaphylactoid reaction, Anaphylactic shock, Anaphylactoid shock.

Nervous system disorders

Very rare: Falcid Paralysis

Cardiac disorders

Uncommon/rare: Tachycardia

Unknown frequency: Kounis syndrome

Vascular disorders

Uncommon/rare: Hypotension

Very rare

- Circulatory collapse and shock

- Flushing

Respiratory, thoracic, and mediastinal disorders

Very rare: Bronchospasm

Skin and subcutaneous tissue disorders

Very rare: Angioneurotic edema, Rash, Urticaria, Erythematous rash

Musculoskeletal disorders

Very rare: muscular weakness *, steroid myopathy *

*after long-term use in the ICU

General disorders and administration site conditions

Uncommon/rare:

Injection site pain/reaction

Drug ineffective

Drug effect/therapeutic response decreased

Drug effect/therapeutic response increased

Very rare:

Face oedema

Malignant hyperthermia

Injury, poisoning and procedural complication

Uncommon/rare:

Prolonged neuromuscular block

- Delayed recovery from anaesthesia

Very rare:

Airway complication of anaesthesia

Additional information on adverse reactions :

Anaphylactic reaction : Although very rare, severe anaphylactic reactions to neuromuscular blocking agents, including rocuronium bromide, have been reported.

Anaphylactic/anaphylactoid reactions are : bronchospasm, cardiovascular changes (e.g. hypotension, tachycardia, circulatory collapse - shock), and cutaneous changes (e.g. angioedema, urticaria). These reactions have, in some cases, been fatal. Due to the possible severity of these reactions, one

should always assume they may occur and take the necessary precautions.

Since neuromuscular blocking agents are known to be capable of inducing histamine release both locally at the site of injection and systemically, the possible occurrence of itching and erythematous reaction at the site of injection and/or generalised histaminoid (anaphylactoid) reactions (see also under anaphylactic reactions above) should always be taken into consideration when administering these drugs.

Prolonged neuromuscular block : The most frequent adverse reaction to non-depolarizing blocking agents as a class consists of an extension of the drug's pharmacological action beyond the time period needed. This may vary from skeletal muscle weakness to profound and prolonged skeletal muscle paralysis resulting in respiratory insufficiency or apnoea.

Myopathy : Myopathy has been reported after the use of various neuromuscular blocking agents in the ICU in combination with corticosteroids.

Local injection site reactions : During rapid sequence induction of anaesthesia, pain on injection has been reported, especially when the patient has not yet completely lost consciousness and particularly when propofol is used as the induction agent.

SYMPTOMS AND TREATMENT OF OVERDOSE :

In the event of overdosage and prolonged neuromuscular block, the patient should continue to receive ventilatory support and sedation. There are two options for the reversal of neuromuscular block : (1) In adults, sugammadex can be used for reversal of intense (profound) and deep block. The dose of sugammadex to be administered depends on the level of neuromuscular block. (2) An acetylcholinesterase inhibitor (e.g. neostigmine, edrophonium, pyridostigmine) or sugammadex can be used once spontaneous recovery starts and should be administered in adequate doses. When administration of an acetylcholinesterase inhibiting agent fails to reverse the neuromuscular effects of rocuronium, ventilation must be continued until spontaneous breathing is restored. Repeated dosage of an acetylcholinesterase inhibitor can be dangerous.

In animal studies, severe depression of cardiovascular function, ultimately leading to cardiac collapse did not occur until a cumulative dose of 750 x ED₅₀ (135 mg/kg rocuronium bromide) was administered.

PHARMACOLOGICAL PROPERTIES :

Pharmacodynamic properties

Mechanism of Action : Rocutroy (rocuronium bromide) is a fast onset, intermediate acting non-depolarising neuromuscular blocking agent, possessing all of the characteristic pharmacological actions of this class of drugs (curariform). It acts by competing for nicotinic cholinceptors at the motor end-plate. This action is antagonised by acetylcholinesterase inhibitors such as neostigmine, edrophonium and pyridostigmine.

Pharmacodynamic effects : The ED₅₀ (dose required to produce 90% depression of the twitch response of the thumb to stimulation of the ulnar nerve) during intravenous anaesthesia is approximately 0.3 mg/kg rocuronium bromide. The ED₅₀ in infants is lower than in adults and children (0.25, 0.35 and 0.40 mg/kg respectively). The clinical duration (the duration until spontaneous recovery to 25% of control twitch height) with 0.6 mg/kg rocuronium bromide is 30-40 minutes. The total duration (time until spontaneous recovery to 90% of control twitch height) is 50 minutes. The mean time of spontaneous recovery of twitch response from 25 to 75% (recovery index) after a bolus dose of 0.6 mg/kg rocuronium bromide is 14 minutes. With lower dosages of 0.3-0.45 mg/kg rocuronium bromide (1-1½ x ED₅₀), onset of action is slower and duration of action is shorter. With high doses of 2 mg/kg, clinical duration is 110 minutes.

Intubation during routine anaesthesia : Within 60 seconds following intravenous administration of a dose of 0.6 mg/kg rocuronium bromide (2 x ED₅₀ under intravenous anaesthesia), adequate intubation conditions can be achieved in nearly all patients of which in 80% intubation conditions are rated excellent. General muscle paralysis adequate for any type of procedure is established within 2 minutes. After administration of 0.45 mg/kg rocuronium bromide, acceptable intubation conditions are present after 90 seconds.

Rapid Sequence Induction : During rapid sequence induction of anaesthesia under propofol or fentanyl/thiopental anaesthesia, adequate intubation conditions are achieved within 60 seconds in 93% and 96% of the patients respectively, following a dose of 1.0 mg/kg rocuronium bromide. Of these, 70% are rated excellent. The clinical duration with this dose approaches 1 hour, at which time the neuromuscular block can be safely reversed. Following a dose of 0.6 mg/kg rocuronium bromide, adequate intubation conditions are achieved within 60 seconds in 81% and 75% of the patients during a rapid

sequence induction technique with propofol or fentanyl/thiopental, respectively.

Paediatric population : Mean onset time in infants, toddlers and children at an intubation dose of 0.6 mg/kg is slightly shorter than in adults. Comparison within paediatric age groups showed that the mean onset time in neonates and adolescents (1.0 min.) is slightly longer than in infants, toddlers and children (0.4, 0.6 and 0.8 min., respectively). The duration of relaxation and the time to recovery tend to be shorter in children compared to infants and adults. Comparing within paediatric age groups demonstrated that mean time to reappearance of T₁ was prolonged in neonates and infants (56.7 and 60.7 min., respectively) when compared to toddlers, children and adolescents (45.4, 37.6 and 42.9 min., respectively).

Geriatric patients and patients with hepatic and/or biliary tract disease and/or renal failure : The duration of action of maintenance doses of 0.15 mg/kg rocuronium bromide might be somewhat longer under enflurane and isoflurane anaesthesia in geriatric patients and in patients with hepatic and/or renal disease (approximately 20 minutes) than in patients without impairment of excretory organ functions under intravenous anaesthesia (approximately 13 minutes). No accumulation of effect (progressive increase in duration of action) with repetitive maintenance dosing at the recommended level has been observed.

Intensive Care Unit : Following continuous infusion in the Intensive Care Unit, the time to recovery of the train of four ratio to 0.7 depends on the level of block at the end of the infusion. After a continuous infusion for 20 hours or more the median (range) time between return of T₁ to train of four stimulation and recovery of the train of four ratio to 0.7 approximates 1.5 (1-5) hours in patients without multiple organ failure and 4 (1-25) hours in patients with multiple organ failure.

Cardiovascular surgery : In patients scheduled for cardiovascular surgery the most common cardiovascular changes during the onset of maximum block following 0.6-0.9 mg/kg rocuronium bromide are a slight and clinically insignificant increase in heart rate up to 9% and an increase in mean arterial blood pressure up to 16% from the control values.

Reversal of muscle relaxation : Administration of acetylcholinesterase inhibitors, (neostigmine, pyridostigmine or edrophonium) at reappearance of T₁ or at the first signs of clinical recovery, antagonises the action of Rocutroy.

Pharmacokinetic properties

Distribution and elimination : After intravenous administration of a single bolus dose of rocuronium bromide, the time course of the plasma concentration runs in three exponential phases. In normal adults, the mean (95%CI) elimination half-life is 73 (66-80) minutes, the (apparent) volume of distribution at steady state conditions is 203 (193-214) ml/kg and the plasma clearance is 3.7 (3.5-3.9) ml/kg/min.

The plasma clearance in geriatric patients and in patients with renal dysfunction is slightly reduced compared to younger patients with normal renal function. In patients with hepatic diseases, the mean elimination half-life is prolonged by 30 minutes and the mean plasma clearance is reduced by 1 ml/kg/min.

When administered as a continuous infusion to facilitate mechanical ventilation for a time period of 20 hours or more, the mean elimination half-life and the mean (apparent) volume of distribution at steady state are increased. A high variability between patients was found in controlled clinical studies, related to the nature and extent of (multiple) organ failure and individual patient characteristics. In patients with multiple organ failure a mean (±SD) elimination half-life of 21.5 (±3.3) hours, an (apparent) volume of distribution at steady state of 1.5 (±0.8) l/kg-1 and a plasma clearance of 2.1 (±0.8) ml/kg/min were found.

Rocuronium bromide is excreted in urine and bile. Excretion in urine approaches 40% within 12 - 24 hours. After injection of a radiolabeled dose of rocuronium bromide, excretion of the radiolabel is on average 47% in urine and 43% in faeces after 9 days. Approximately 50% is recovered as rocuronium bromide.

Biotransformation : No metabolites are detected in the plasma.

SPECIAL PRECAUTION FOR USE AND OTHER HANDLING :

Compatibility studies with the following infusion fluids have been performed: In normal concentrations of 0.5mg/ml and 2.0 mg/ml Rocutroy has been shown to be compatible with 0.9% sodium chloride, 5% dextrose, 5% dextrose in 0.9% sodium chloride, sterile water for injection, Haemacel and Lactated Ringers. Administration should be begun immediately. After mixing and should be completed within 24 hours. Any unused medicinal product or waste should be disposed of in accordance to local requirements.

INCOMPATIBILITIES :

Physical incompatibility has been documented for rocuronium when added to solutions containing the following drugs: amphotericin, amoxicillin, azathioprine, cefazolin, cloxacillin, dexamethasone, diazepam, enoximone, erythromycin, famotidine, furosemide, hydrocortisone sodium succinate, insulin, Intralipid, methohexital, methylprednisolone, prednisolone sodium succinate, thiopental, trimethoprim and vancomycin.

Rocuronium must not be mixed with other medicinal products except those mentioned in SPECIAL PRECAUTIONS FOR USE AND OTHER HANDLING. If Rocuronium is administered via the same infusion line that is also used for other drugs, it is important that this infusion line is adequately flushed (e.g. with 0.9% NaCl) between administration of rocuronium and drugs for which incompatibility with rocuronium has been demonstrated or for which compatibility with rocuronium has not been established.

EXCIPIENTS:

Sodium Chloride, Sodium Acetate (Trihydrate), Glacial Acetic Acid, Water for Injection.

STORAGE CONDITION :

Unopened Vials : Store between 2° C to 8° C. Protect from light.

After dilution : After dilution with infusion fluids, (see SPECIAL PRECAUTIONS FOR USE AND OTHER HANDLING) chemical and physical in-use stability has been demonstrated for 72 hours at 30°C.

From a microbiological point of view, the diluted product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user/administrator.

SHELF LIFE :

Unopened Vials : 24 months, provided it is stored under the prescribed conditions. See STORAGE CONDITION.

Opened Vials : Since Rocutroy does not contain a preservative, the solution should be used immediately after opening the vial.

After dilution

After dilution with infusion fluids, chemical and physical in-use stability has been demonstrated for 72 hours at 30°C.

From a microbiological point of view, the diluted product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user/administrator.

PRESENTATION :

ROCUTROY Rocuronium Bromide 10mg/ml Solution for Injection or Infusion is filled in a 5mL Flint tubular clear glass vial USP Type-I with 20 mm grey coloured bromo butyl rubber stopper and 20mm light pink flip off seal packed in printed carton with printed insert.

DATE OF REVISION :

30/06/2020

PRODUCT REGISTRATION HOLDER / IMPORTED BY :

Unimed Sdn. Bhd.

No. 53, Jalan Tembaga SD5/2B,
Bandar Sri Damansara,
52200 Kuala Lumpur, Malaysia.

Manufactured by :

TroiKaa

TroiKaa Pharmaceuticals Ltd.

C-1, Sara Industrial Estate, Selaqui, Dehradun-248197, Uttarakhand, (India)

DJRT.MAL.EU