

PACKAGE INSERT

QILU PHARMACEUTICAL (INNER MONGOLIA) 15% W/W BACITRACIN ZINC PREMIX

Brand of Product Name	Qilu Pharmaceutical (Inner Mongolia) 15% w/w Bacitracin Zinc Premix
Strength of Active Substance(s)	Each kilogram product contains 150 g bacitracin zinc
Product Description	Slightly brown or tawny granule
Pharmacodynamics :	Bacitracin is produced by <i>Bacillus licheniformis</i> and <i>Bacillus subtilis</i>, and is a mixture of several closely related polypeptides, mainly consisting of bacitracin A, B1, B2, C and a small portion of bacitracin F (with relative ratio's of approximately 53%, 22%, 13%, 8%, and 4%, respectively, although this may vary according to producing species and/or production conditions). The zinc salt adds stability particularly during storage of the product. Bacitracin is mainly active against Gram-positive microorganisms. The activities of each of bacitracins B1, B2, and C constitute about 90% of the antibacterial activity of bacitracin A. Bacitracin F is almost inactive.
Pharmacokinetics :	After oral application to rats, chickens and pigs, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In rats, chickens and pigs, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolized to amino acids and smaller peptides via the main metabolite desamidobacitracin, which is microbiologically inactive. Main metabolites in faeces are bacitracin A, B1, B2, F, desamidobacitracin and catabolic peptides. In urine and bile only hydrolytic cleavage products (di- and tripeptides) are present.
Indication	For the treatment of Necrotic Enteritis (<i>Clostridium Perfringens</i> Types A and C) and as an aid in the prevention of Necrotic Enteritis in poultry.
Dosage	Treatment of Necrotic Enteritis: Add 1.33kg of Bacitracin Zinc Premix 15% w/w Granule per tonne of feed to produce a concentration of 200g of bacitracin per tonne of finished feed. Feed for 5-7 days. Aid in the Prevention of Necrotic Enteritis: Add 267g of Bacitracin Zinc Premix 15% w/w Granule per tonne of feed to produce a finished feed (corresponding to 40g of bacitracin per tonne of feed). Medicated feed should be fed to poultry during the period when they are at most risk of necrotic enteritis, usually between 2 and 12 weeks. Bacitracin Zinc Premix 15% w/w Granule must be thoroughly mixed in the feed to assure even distribution. It is recommended that the Bacitracin Zinc Premix 15% w/w Granule is premixed with 10-20 kg of feed before adding to the bulk feed.

Manufactured By:
Qilu Pharmaceutical (Inner
Mongolia) Co. Ltd
Address:

No.2 Wei Si Road, Jinchuannan District,
Economy & Technology Development
Zone, Hohhot City, China (010080)

Mode of Administration	Oral administration by mixing Bacitracin Zinc Premix 15% with feed.
Contraindication	Not known.
Warning and Precaution	Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period. Safety Directions: Will irritate the eyes, nose and throat. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into feed, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles and disposable face mask covering mouth and nose. Keep out of reach of children/ Jauhi ubat daripada kanak-kanak.
Interaction with other medical product:	None known.
Pregnancy and lactation	It is prohibited to be used for breeding poultry and breeding livestock.
Target species	Poultry
Adverse Effect	None known.
Symptoms and treatment of overdose	Not known.
Withdrawal period/s	Meat: Zero (0) days Eggs: Zero (0) days
Storage conditions	Storage below 25°C (Air conditioning) in a dry place. Protect from light.
Dosage forms and packaging available	25 kg
Name and Address of Manufacture/Marketing authorization holder	Gladron Chemicals Sdn. Bhd No. 7, Jalan TP 7, UEP Industrial Park, 40400 Shah Alam, Selangor D.E., Malaysia
Date of Revision of Package Insert	May 2020

Distributed by:
Gladron Chemicals Sdn. Bhd
Address: No. 7, Jalan TP 7, UEP
Industrial Park, 40400 Shah Alam,
Selangor D.E., Malaysia.