

PACKAGE LEAFLET FOR Chanazine 10 % w/v Solution for Injection.

NAME OF THE VETERINARY MEDICINAL PRODUCT

Chanazine 10% w/v Solution for Injection.

COMPOSITION

Each ml contains 100 mg Xylazine Base

As preservatives:

Methyl Parahydroxybenzoate 0.18% w/v and Propyl Parahydroxybenzoate 0.02% w/v.

DESCRIPTION:

Clear colourless solution, free from particulate matters

PHARMACODYNAMICS:

Xylazine is an α_2 -adrenergic drug with sedative, analgesic and muscle relaxing properties which acts via the CNS. Xylazine is thought to act by activation of the central presynaptic α_2 -receptors. Activation of these central α_2 -receptors seems to regulate central dopamine and norepinephrine storage or release. Xylazine's analgesic and sedative actions are related to its central nervous system depression, while the muscle relaxant effects are due to the inhibition of the intraneural transmission of impulses in the central nervous system.

PHARMACOKINETICS:

Onset and duration of sedation is species-dependent varying from 5 to 15 minutes in horses

TARGET SPECIES: Horses

INDICATIONS:

Indications for use, specifying the target species

Xylazine is a sedative with analgesic and muscle relaxant properties. The product is for use in horses only, in cases where sedation is required including:

1. Handling fractious animals e.g. for transportation.
2. Medical examinations e.g. X-ray examinations, removal of bandages; examination of the penis and oral cavity.
3. Premedication for minor superficial operations, and local or regional anaesthesia.
4. Elimination of defaecation when examining and treating the vagina, uterus and hindquarters.

RECOMMENDED DOSAGE:

The product is given by slow intravenous injection. Dosage depends on the degree of sedation required and the response of the animal, and is 0.5 – 1 ml/100 kg (0.5 - 1 mg/kg bodyweight). Nervous or excitable horses may require higher doses. Older horses and those having undergone severe physical exertion before treatment should receive the lowest dose rate.

Animals do not usually become recumbent after administration and light to deep sedation with variable degree of analgesia is obtained. Effects are usually seen within 5 minutes and persist for approximately 20 minutes. The product may be employed as a premedication to barbiturate anaesthesia or in combination with regional or local anaesthesia.

MODE OF ADMINISTRATION: INTRAVENOUS

CONTRA-INDICATION/SAFETY DATA:

Do not administer by the intra-carotid route.

Careful consideration should be given before administering to animals exposed to stress conditions such as extreme heat, cold, high altitude or fatigue.

WARNINGS AND PRECAUTIONS

Special warnings for each target species

None known

Special precautions for use

i. Special precautions for use in animals

Should be used with caution when pulmonary disease is suspected.

Do not exceed the recommended dosage

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Horses sedated with xylazine usually remain standing and may still kick with accuracy.

Precaution should be taken to avoid accidental injection / self-injection.

In the case of accidental oral intake or self-injection, seek medical advice immediately and show the package leaflet to the doctor but DO NOT DRIVE as sedation and changes in blood pressure may occur.

Avoid skin, eye or mucosal contact.

Immediately after exposure, wash the exposed skin with large amounts of fresh water.

Remove contaminated clothes that are in direct contact with skin.

In the case of accidental contact of the product with eyes, rinse with large amounts of fresh water. If symptoms occur, seek the advice of a doctor.

If pregnant women handle the product, special caution should be observed not to self inject as uterine contractions and decreased foetal blood pressure may occur after accidental systemic exposure.

Advice to Doctors: Xylazine is an alpha2-adrenoreceptor agonist. Symptoms after absorption may involve clinical effects including dose-dependent sedation, respiratory depression, brachycardia, hypotension, a dry mouth, and hyperglycaemia. Ventricular arrhythmias have also been reported. Respiratory and haemodynamic symptoms should be treated symptomatically

INTERACTIONS WITH OTHER MEDICATIONS

None known.

INCOMPATIBILITIES:

None known.

PREGNANCY AND LACTATION

Xylazine should not be administered during the later stages of pregnancy because of the risk of inducing premature parturition.

As the safety of xylazine use during organogenesis has not been fully demonstrated by current methods it should be used with caution during the first month of pregnancy

SIDE EFFECTS:

Side effects of bradycardia, cardiac arrhythmia and polyuria may occur. Following intravenous administration a transient rise followed by a fall in blood pressure usually occurs.

SYMPTOMS AND TREATMENT OF OVERDOSE

Sedation is dose dependent and the occurrence of side effects will increase and worsen at higher dosage than recommended.

WITHDRAWAL TIMES:

Animals must not be slaughtered for human consumption during treatment.

Horse:	Not to be used in horses intended for human consumption. Treated horses may never be slaughtered for human consumption..

PACKING : 50 ml per vial.10 vial in a carton box

STORAGE:

Store below 25 0C
Protect from light and moisture

PROPOSED SHELF LIFE:

3 years as packaged for sale
DISCARD AFTER 24 HOURS OF OPENING

WARNING: FOR VETERINARY USE ONLY
KEEP OUT OF THE REACH OF CHILDREN
JAUHKAN DARI KANAK-KANAK

PRODUCT REGISTRATION HOLDER :

Km Vet Pharm Sdn Bhd
No.1, Jln Industri Batu Caves 1/1,
Taman Perindustrian Batu Caves 68100
Batu Caves, Selangor, Malaysia

MANUFACTURER:

DIVASA-FARMAVICS.A. ,
(DFV), Ctra. Sant Hipolit ,
km 71,08503 Gurb –Barcelona, Spain

Product Owner:

Chanelle Pharmaceuticals Manufacturing Limited
Dublin Road, Loughrea, County Galway
Ireland

DATE OF REVISION: FEBRUARY 2020