

|                                              |                                    |            |                            |
|----------------------------------------------|------------------------------------|------------|----------------------------|
| Product:                                     | Esberitox (family)                 | Fontsize:  | 24 pt; 8 pt                |
| Country:                                     | MALAYSIA (MAL)                     | Used Font: | Optima<br>Frutiger LT Std. |
| Packaging:                                   | patient information leaflet        | Colours:   | Black                      |
| Manufacturer:                                | Schaper & Brümmer GmbH &<br>Co. KG |            |                            |
| Drawing No:                                  | 0000/049/00                        | 1. proof:  | 17.04.2019                 |
| Pharmacode:                                  | 00000                              | 2. proof:  | 24.04.2019                 |
| Dimension:                                   | 148 x 210 mm                       | 3. proof:  | 12.12.2019                 |
| Page:                                        | 1 of 2                             | 4. proof:  | 13.12.2019                 |
|                                              |                                    | 5. proof:  | 19.12.2019                 |
| <input type="checkbox"/> Correction required |                                    | 6. proof:  | 19.12.2019                 |
| <input type="checkbox"/> Final Version/Date: |                                    | 7. proof:  | 11.02.2020                 |
| <input type="checkbox"/> Approval            |                                    |            |                            |

Package leaflet: Information for the user

Esberitox®  
family

for application in adults and children from the age of 2 years  
dry extract from a mixture of wild indigo rootstock, purple coneflower root,  
pale coneflower root and thuja herba

**Read all of this leaflet carefully before you start taking this medicine  
because it contains important information for you.**

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to your doctor or pharmacist if you do not feel better or if you feel worse after 7 days.

What is in this leaflet

1. What Esberitox®family is and what it is used for
2. What you need to know before you take Esberitox®family
3. How to take Esberitox®family
4. Possible side effects
5. How to store Esberitox®family
6. Contents of the pack and other information

1. What Esberitox®family is and what it is used for

Esberitox®family is traditionally used for the relief of cold. This is a traditional medicine.

2. What you need to know before you take Esberitox®family

2.1 Do not take Esberitox®family

- if you are allergic to the pharmacologically active ingredients (wild indigo rootstock, purple coneflower root, coneflower root, thuja herba), or any of the other ingredients of this medicine, or other plants of the Asteraceae (comPOSITE) family.

2.2 Warnings and precautions

The use is not recommended in cases of progressive systemic disorders, autoimmune diseases, immunodeficiencies, immunosuppression and diseases of the white blood cell system.

2.3 Children

This medicine is not suitable for children under the age of 2 years.

2.4 Other medicines and Esberitox®family

Interactions are not known.

Please, nevertheless inform your doctor or pharmacist, if you take or have recently taken other medicines, including medicines obtained without a prescription.

2.5 Pregnancy and breast-feeding

Contraindicated in pregnant women. Insufficient reliable data in breastfeeding women.

2.6 Driving and using machines

No special precautions are required.

3. How to take Esberitox®family

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

3.1 Unless otherwise prescribed by your doctor, the recommended dose is

Adults: Take **3 tablets 3 times daily** after meal.  
Children 7 to 12 years: Take **2 tablets 3 times daily** after meal.

|                                              |                                 |            |                            |
|----------------------------------------------|---------------------------------|------------|----------------------------|
| Product:                                     | Esberitox (family)              | Fontsize:  | 24 pt; 8 pt                |
| Country:                                     | MALAYSIA (MAL)                  | Used Font: | Optima<br>Frutiger LT Std. |
| Packaging:                                   | patient information leaflet     | Colours:   | Black                      |
| Manufacturer:                                | Schaper & Brümmer GmbH & Co. KG |            |                            |
| Drawing No:                                  | 0000/049/00                     | 1. proof:  | 17.04.2019                 |
| Pharmacode:                                  | 00000                           | 2. proof:  | 24.04.2019                 |
| Dimension:                                   | 148 x 210 mm                    | 3. proof:  | 12.12.2019                 |
| Page:                                        | 2 of 2                          | 4. proof:  | 13.12.2019                 |
|                                              |                                 | 5. proof:  | 19.12.2019                 |
| <input type="checkbox"/> Correction required |                                 | 6. proof:  | 19.12.2019                 |
| <input type="checkbox"/> Final Version/Date: |                                 | 7. proof:  | 11.02.2020                 |
| <input type="checkbox"/> Approval            |                                 |            |                            |

Children 2 to 6 years: Take **1 – 2 tablet(s)** **3 times daily** after meal.

Please take Esberitox®family in the morning, in the middle of the day and in the evening with sufficient liquid, preferably 1 glass of water. The tablets may also be chewed.

**3.2 Duration of use**

Start the treatment as early as possible after the first symptoms appear and continue taking Esberitox®family until the symptoms disappear. However, take note of the details on precautionary measures of section 2.2 and the details on side effects of section 4 in this package leaflet.

Do not take Esberitox®family longer than 10 days without medical advice.

**3.3 If you take more Esberitox®family than you should**

Toxications with Esberitox®family have not been known until now. If you inadvertently have taken one or two tablets more than you should, this has normally no adverse effects. If you have taken a considerably larger quantity of Esberitox®family than you should, please consult a doctor. He/she can decide on required measures to be taken, if necessary.

**3.4 If you forget to take Esberitox®family**

Do not take a double dose, but continue the intake as described in this package leaflet or as prescribed by your doctor.

**4. Possible side effects**

- Hypersensitivity reactions, e. g. skin rash, itching, facial swelling, shortness of breath, blood pressure decrease
- Abdominal pain, nausea, diarrhea
- Dizziness

**5. How to store Esberitox®family**

Keep below 30° Celsius. Protect from light and moisture.

Keep away from children. Jauhkan dari kanak-kanak.

**6. Contents of the pack and other information**

**What Esberitox®family contains**

The active substance is:  
Each 190 mg tablet contains:  
3.2 mg extract (as dry extract) from Baptisia tinctoria L. (wild indigo), radix corresponding to approx. 11 mg wild indigo rootstock;  
Echinacea purpurea Moench (purple cone-flower), radix corresponding to approx. 4 mg dried root from purple coneflower;  
Echinacea pallida Nutt. (pale coneflower), radix corresponding to approx. 4 mg dried root from pale coneflower  
and Thuja occidentalis L. (thuja), herba corresponding to approx. 2 mg dried herba from thuja.

The other ingredients are:  
Betadex, mannitol (Ph. Eur.), glycerol dibehenate (Ph. Eur.), magnesium stearate (Ph. Eur.) [herbal], saccharin sodium, citric acid monohydrate, lemon flavoring (natural).

**What Esberitox®family looks like and contents of the pack**

Round, marbled, beige-coloured tablets

Esberitox®family is available in packs with 100 tablets.

**Product Registration Holder**

Quanstar Biotech Sdn Bhd, B-10-2,  
Northpoint Office  
Mid Valley City, No. 1, Medan Syed Putra  
Utara,  
59200 Kuala Lumpur  
Malaysia

**Manufacturer**

 Schaper & Brümmer GmbH & Co. KG  
Bahnhofstr. 35  
38259 Salzgitter  
Lower Saxony,  
Germany  
Tel. +49 53 41 3 07-0  
Fax +49 53 41 3 07-124  
info@schaper-bruemmer.de

**This package information leaflet was last revised in February 2020**