

Solution for injection

Vime Tylovet

50mg/ml

COMPOSITION:

Each ml contains:
Tylosin 50 mg
With 4.0% (v/v) Benzyl Alcohol as a preservative

DESCRIPTION

The finished product is yellow transparent solution.

TARGET SPECIES

Cattle (Beef, calves, dairy, non-lactating), swine

PHARMACODYNAMICS

Tylosin is a macrolide antibiotic and exerts its antibiotic activity by a similar mechanism to other macrolides i.e. binding to the 50S fraction of the ribosomes resulting in an inhibition of the synthesis of proteins. Tylosin has mainly a bacteriostatic activity. Tylosin has an antibiotic effect against Gram positive cocci (Staphylococci, Streptococci), Gram positive bacilli (*Erysipelothrix* spp.), some Gram-negative bacilli (*Pasteurella* spp., *Mannheimia* spp.) and *Mycoplasma* spp. Resistance to macrolides is usually plasmid-mediated but modification of ribosomes may occur through chromosomal mutation. Resistance can occur by i) decreased entry into bacteria (most common with the gram-negative bacteria), ii) synthesis of bacterial enzymes that hydrolyse the drug and, iii) modification of the target (the ribosome). This latter resistance type may also lead to cross-resistance with other antibiotics that preferentially bind to bacterial ribosome. Gram-negative anaerobic bacteria are often resistant.

Resistance of *Brachyspira hyodysenteriae* to tylosin has been reported.

PHARMACOKINETICS

Absorption: Following intramuscular injection, tylosin blood levels peak 1-2 hours post-injection. Duration of activity is approximately 12 hours.

Distribution, Biotransformation and Elimination: Tylosin levels of 1.4 to 1.6 and 2.2 to 6.7 µg/ml were recorded in serum and lung tissue respectively following intramuscular injection of 8.8 mg/kg bodyweight in pigs. Measurable amounts of tylosin were still present in both serum and lung tissue at 12 hours post injection.

Tylosin concentrations were greater in lung tissue than serum at all sample times.

INDICATION

Swine:

Indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Cattle (Beef and Non-lactating Dairy):

For use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

ROUTE OF ADMINISTRATION

Intramuscular route

RECOMMENDED DOSE

Swine:

9 mg per kg of body weight (1 ml per 5.6 kg) twice daily; not to exceed 3 days.

Cattle (Beef and non-lactating dairy):

18 mg per kg of body weight one time daily (1 ml per 2.8 kg); not to exceed 5 days.

CONTRA-INDICATION

Do not administer to chickens or turkeys in which intramuscular injection may be fatal.

Do not administer to horses or other equines in which injection of tylosin may be fatal.

Do not administer to animals with known hypersensitivity to tylosin, other macrolides or any of the excipients.

INTERACTION WITH OTHER MEDICAL PRODUCT

Lincosamide and aminoglycoside antibiotics can antagonise the action of tylosin.

WARNINGS AND PRECAUTIONS

Do not mix this product with other injectable solutions as this may cause a precipitation of the active ingredients.

Do not attempt injection into pigs weighing less than 2.8 kg (0.5 ml) unless the syringe is capable of accurately delivering 0.1 ml.

Do not administer to horses or other equines. Injection of tylosin in equines has been fatal.

- Keep out of reach of children
- Jauhkan daripada kanak-kanak

PREGNANCY AND LACTATION

Studies in laboratory animals have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species.

Use only in accordance with the benefit/risk assessment by the responsible veterinarian.

SIDE EFFECTS

Adverse reactions, including shock and death, may result from overdosage in baby pigs.

SYMPTOMS AND TREATMENT OF OVERDOSE

Pigs and calves: Intramuscular injection of 30 mg/kg bodyweight per day for five days produced no adverse effects.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

STORAGE CONDITION

Store in a dry place not exceeding 30°C. Protect from light.

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 24 hours

WITHDRAWAL PERIOD

Cattle (Meat): 21 days.

Pigs (Meat): 14 days.

A withdrawal period has not been established for this product in pre-ruminating calves.

PACKAGING: 100 ml

REVISED DATE: Jul 03, 2020