PRODUCT INFO

SLID #	243462AF	Product Name:	Surmax 100 Premix
Item Code:	BG243462X	Component:	Printed Bag
Product Code:	AF0375	Pack Size:	25kg
Previous Item Code:	BG210232X		

ARTWORK INFO

Template:	BG_41Gx94Gx100mm_Avilo_v4	Packaging Spec(s):	BG0103
Barcodes/Type:	N/A	Web Info:	N/A
		Minimum Core Data Point Size:	24 pt
OTIN:	N/A	Proof #:	Proof Date: 04-06-OCT-2025

Key Printed Ink: BLACK; PMS 1805

ELANCO ARTWORK LEGEND V10

top of bag

JOINT

LEFT GUSSET

FRONT

RIGHT GUSSET

BACK

Elanco™

AF0375

Surmax™ 100

TM

Net Weight: **25 kg**

For Use in Chicken and Swine Feeds

Granular medicated premix for the preparation of medicated animal feed

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

JAUHI UBAT DARI KANAK-KANAK

ACTIVE INGREDIENT:
100 g per kg of Avilamycin.

INDICATIONS:
1. For the prevention of necrotic enteritis due to *Clostridium perfringens* in growing broiler chickens.
2. For reduction in the incidence and severity of post-weaning diarrhea associated with *Escherichia coli* in pigs. Avilamycin is to be used in pigs that are at risk of developing, but not yet showing clinical signs of, diarrhea in the presence of pathogenic *E. coli*.

Notes:
1. Comprehensive necrotic enteritis management should be planned in consultation with veterinary and nutritional poultry expertise, and based on flock health history and exposure to necrotic enteritis risk factors.
2. Comprehensive management of diarrhea due to *E. coli* should be planned in consultation with a veterinarian and animal nutritionist.
3. As Surmax Premix is indicated for use in weanling pigs, responsible antimicrobial use limits its use to pigs less than 14 weeks of age.

DOSAGE AND ADMINISTRATION:
For Indication 1: Thoroughly mix 0.15 to 0.30 kg of Surmax Premix per 1000 kg of feed to yield 15 to 30 ppm of avilamycin.
For Indication 2: Thoroughly mix 0.80 kg of Surmax Premix per 1000 kg of feed to yield 80 ppm of avilamycin.

IMPORTANT: Must be thoroughly mixed in broiler chicken or swine feeds before use. For better dispersion, it is recommended that Surmax Premix be thoroughly mixed with a small quantity of the feed ingredients (totalling at least 10 kg) prior to the manufacturing of the medicated feed.

CONTRAINDICATION:
None known.

Manufactured by:
Elanco UK AH Limited
Fleming Road, Liverpool
L24 9LN, UK

For:
Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6, Tower Block
The Bousteador, No. 10, Jalan PJU 7/6
Mutiara Damansara, 47800 Petaling Jaya
Selangor, Malaysia

TAKE TIME
OBSERVE LABEL DIRECTIONS

STORE AT OR BELOW 30°C

The product will remain stable in the finished feed for 3 months.

Surmax, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2025 Elanco or its affiliates

BG243462X

Batch No.: Mfg Date: Expiry Date:

Elanco™ Surmax™ 100

Elanco™ Surmax™ 100

Elanco™

AF0375

Surmax™ 100

PRODUCT DESCRIPTION:
Light to medium brown free flowing granular material which may contain light coloured particles.

PHARMACODYNAMICS:
Avilamycin is an orthosomycin antibiotic complex produced by the fermentation of Streptomyces viridochromogenes. Orthosomycin antibiotics are divided into two groups: those that contain an aminoacyclitol residue and those that are esters of dichlorisoeverninic acid. Avilamycin falls into the latter groups as do the evernimicins.

Avilamycin inhibits protein synthesis. It is thought to bind to 50S ribosomal subunit. It prevents the association of IF2 which inhibits the formation of the mature 70S initiation complex and the correct positioning of the tRNA in the aminoacyl site.

PHARMACOKINETICS:
The oral bioavailability of avilamycin in pigs, chickens and rat was low. This limited the capability to quantitatively demonstrate the distribution of avilamycin to tissues. Thus standard pharmacokinetic data were not available.

Most distribution studies were performed in pigs and chickens using radiolabelled avilamycin. A comparative study indicated that in rat the metabolism and distribution was comparable to pigs. Avilamycin is extensively metabolized and exhibits low tissue residue levels when administered orally to rats or swine. Therefore metabolites were only characterized in liver, due to low residue levels in the other tissues. Flambic acid is the primary metabolite in liver (15 to 20%) and faeces (40 to 50%). In liver no microbiologically active residues were detected. In chicken radioactive residues could not be characterized due to low levels of residues in the tissues and excreta. No metabolism data are available for rabbits or turkeys. When administered orally to swine, avilamycin was primarily excreted via the faeces (95%). The bulk of the radioactivity is excreted within the first week. Biliary excretion is of minor importance (estimated to be 7% or orally administered dose).

WARNINGS AND PRECAUTIONS:
In swine, the effectiveness of Surmax has only been demonstrated when avilamycin-medicated feed was offered before the onset of diarrhea. Use of Surmax for the treatment of established cases of clinical diarrhea in weanling pigs has not been demonstrated.

Do not use in feeds containing pellet binding agents with the exception of calcium lignosulfate pellet binder.

Safety Directions:
When handling the product, avoid inhalation, oral exposure, and direct contact with skin or eyes. Use protective clothing, impervious gloves, goggles, and an approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention.

WITHDRAWAL PERIOD:
No withdrawal period is required for broiler chickens or swine when treated according to the label.

Do not use in laying hens.

INTERACTIONS WITH OTHER MEDICINAL PRODUCTS:
None.

PREGNANCY AND LACTATION:
In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.

ADVERSE EFFECTS:
If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

SYMPTOMS OF OVERDOSE AND TREATMENT:
Stop medication and to seek for veterinarian advice if there is sign of overdosage.

DISPOSAL OF CONTAINER:
No special package disposal is required.

CONTROLLED MEDICINE
MAL20096022HA
Revised Date: 11 September 2025

BG243462X

Area designated for UN Number – applied by Print Vendor

base of bag

120mm

+10

940. mm

1000. mm

50. mm

36. mm

36. mm

50. mm

6

5

8+

50. mm

100. mm

410. mm

100. mm

360. mm