
LYSOME SYRUP

Reg. No.: MAL20101996XE

Description:

Clear, light yellow liquid.

Composition:

Each 5 ml syrup contains:	Lysozyme Hydrochloride	25.0 mg
	Potassium Sorbate (as preservative)	1.9 mg
	Sodium Benzoate (as preservative)	0.12 mg

Therapeutic Properties:

LYSOME Syrup is a preparation containing a purified Lysozyme Hydrochloride derived from egg white.

Lysozyme is a basic protein composed of 129 amino acids of 18 different types with a molecular weight of about 14,300. Lysozyme Hydrochloride occurs as a white, crystalline powder. It has no odour and has a sweet taste. It is very soluble in water.

Lysozyme is widely distributed in tears, nasal discharge, saliva, etc. in the case of humans. It has been discovered to possess anti-inflammatory, tissue repairing, mucopolysaccharidase and anti-hemorrhagic actions, etc. Thus, Lysozyme which has various actions plays an important role in the living body as an enzyme responsible for the physiological protective mechanism.

Clinically, Lysozyme has been used in expectation of its anti-inflammatory, expectorant and anti-hemorrhagic effects, etc. in many fields.

Indications:

Chronic paranasal sinusitis.

Difficulty of expectoration caused by respiratory diseases.

Haemorrhage during and after operations (dental, urological fields)

Contraindications:

Contraindicated in patients who have shown hypersensitivity to Lysozyme or patients whose family members have a history of allergic reactions to Lysozyme.

Side Effects / Adverse Reactions:

1) Shock

Since Lysozyme may cause shock symptoms, patients taking Lysozyme should be observed closely. If pallor facial, coldness of limbs, blood pressure dropped, cyanosis or unconsciousness occurred, discontinue the administration and appropriate treatment should be given.

2) Hypersensitivity

May cause hypersensitivity reactions, such as eruption or redness. Discontinue the medication if these symptoms occur.

3) Digestive System

May cause diarrhea, anorexia, discomfort in the stomach, nausea or vomiting, and rarely cause stomatitis.

Dosage and Administration:

Usual daily doses are:

Less than 2 years old : 1 - 2 ml three times per day

2 - 6 years old : 2 – 2.5 ml three times per day

7 - 14 years old : 2.5 - 4 ml three times per day

Usage in Pregnancy:

As safety of Lysozyme has not been established, use with caution in pregnant women or women suspected of being pregnant.

Drug Interactions:

None known

Overdosage and Treatment:

No cases of acute overdose have been reported with Lysozyme Hydrochloride. The oral LD50 of Lysozyme Hydrochloride is greater than 20 g/kg in mice. Early gastric lavage may be indicated depending on the amount of overdosage and nature of symptoms.

Incompatibilities:

None Known

Storage Conditions:

Store below 25°C. Keep out of reach of children.

Shelf-life:

3 years

Pack Size:

In PE bottle of 100 ml and 500 ml.

Unison Nutraceuticals Sdn Bhd

Manufacturer:

UNISON NUTRACEUTICALS SDN BHD

No. 13, Jalan TTU 52, Taman Tasik Utama, Ayer Keroh, 75450 Melaka, MALAYSIA

Distributed By:

WELLCHEM PHARMACEUTICALS PTE. LTD.

221 Henderson Road, #04-15, Henderson B Building, SINGAPORE 159557

A product of Nippon Shinyaku Co. Ltd., Kyoto, JAPAN

The information in this leaflet is limited. If in doubt, please consult your physician or pharmacist.