



WOUND OINTMENT

For external use only

Please read these instructions before use. Use this medicine only as directed by a physician.

Product Name: MEBO® Wound Ointment

Each 1g ingredient contains:

Rhizoma Coptis Chinensis extract 44mg, Cortex Phellodendron Chinense extract 44mg, Radix Scutellaria Baicalensis extract 44mg

Product Description: Light to dark brownish-yellow color with sesame oil odor.

Indications: Traditionally used for relief of minor burns and cuts.

Specification: 10 grams per tube; 20 grams per tube; 40 grams per tube

Dosage and Administration:

Intended for external use only. Adult: Apply a thin film (less than 1mm thickness) of MEBO® Wound Ointment to the wound surface every 4 to 6 hours per day. Before reapplying, the residual ointment and liquefied products should be wiped off.

Adverse Reaction: Unknown

Contraindication: This product contains sesame oil and is not suitable for user who is allergic/hypersensitive to sesame oil.

Precaution: Patients developing systemic signs derived from burn wounds should be treated under the supervision of doctors. Extreme heat or repeated compression pressing of the MEBO tube may make the ointment thinner and more fluid like, but this should not hinder or change its effects/efficacy. However, if thinning occurs, immerse the tube with the cap tightly screwed on in water that has been boiled, then let the tube stand vertically with the capside down to allow it to cool to room temperature so that its consistency returns to normal.

Warning:

- i. This is a Traditional Medicine. Please consult your pharmacist/doctor if symptoms persist/worsen;
- ii. Keep out of reach of children/Jauhkan daripada capaian kanak-kanak. For external use only;
- iii. Not to be taken by babies, children under 12 years of age, pregnant women or lactating mothers. Consult your practitioner if you have conditions such as: Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency, Haemolytic anemia, Glaucoma, Diabetes, High Blood Pressure, History of cardiovascular disease and if you are using Paclitaxel, Cyclosporin, or other chemotherapeutic agents;
- iv. Not to be taken by pregnant women or lactating mothers.

Storage Condition: Store below 30°C. Protect from light and moisture.

Shelf Life: 36 months

Product registration holder: Mebo Healthcare Sdn. Bhd.

Address: Lot 4-401&4-402, Level 4, The Starling Mall, No.6, Jalan SS21/37, Damansara Uptown, 47400 Petaling Jaya, Selangor, Malaysia

Manufacturer: Shantou MEBO Pharmaceutical Co., Ltd.

Address: No.13, Southern Yujin Road, Zhujin Industrial District, Shantou City, Guangdong, China