

Veterinary Package Insert

LIN-SPECT 222/445 MG/G WATER SOLUBLE POWDER (25.2% Lincomycin Hydrochloride & 66.3% Spectinomycin Hydrochloride)

PRODUCT DESCRIPTION

White to off-white colour powder, which contains 252g of lincomycin hydrochloride (222g of lincomycin) and 663g of spectinomycin hydrochloride (444.7g of spectinomycin) in 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Pharmacodynamics:

This is a combination of two antibiotics, lincomycin and spectinomycin, having a complementary spectrum of activity.

Lincomycin

Lincomycin is active against gram-positive bacteria, some anaerobic gram-negative bacteria and mycoplasmas. It has little or no action against gram-negative bacteria such as *Escherichia coli*.

Spectinomycin

Spectinomycin is an aminocyclitol antibiotic derived from *Streptomyces spectabilis*, it has bacteriostatic activity and is active against *Mycoplasma* spp. and against some gram-negative bacteria such as *E. coli*.

The mechanism by which spectinomycin administered orally acts on pathogens at the systemic level despite a poor absorption is not fully elucidated, and might rely partly on indirect effects on the gut flora.

Pharmacokinetics:

Lincomycin

In pigs, lincomycin is rapidly absorbed following oral administration. Lincomycin pharmacokinetic studies in pigs show that lincomycin is bioavailable when given intravenously, intramuscularly or orally. The average of the half-lives of elimination of all routes of administration is 2.82 hours in pigs.

In chickens treated with the veterinary medicinal product in drinking water at the target dose of 50 mg/kg body weight of total activity (at a ratio of 1:2 lincomycin:spectinomycin) for seven consecutive days, C_{max} after first offering of medicated water was calculated to be 0.0631µg/ml. C_{max} occurred at 4 hours after introduction of the medicated water.

Spectinomycin

Studies performed in various animal species have demonstrated that spectinomycin undergoes limited absorption from the intestine (less than 4–7%) after oral administration. Spectinomycin exhibits little tendency to protein binding and is poorly liposoluble.

INDICATION

Pigs

For the treatment and metaphylaxis of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*, and associated enteric pathogens (*Escherichia coli*) susceptible to lincomycin and spectinomycin.

The presence of the disease in the group must be established before the product is used.

Chickens

For the treatment and metaphylaxis of chronic respiratory disease (CRD) caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to lincomycin and spectinomycin, and associated with a low mortality rate.

The presence of the disease in the flock must be established before the product is used.

TARGETED SPECIES

Pigs and chickens.

RECOMMENDED DOSAGE AND ADMINISTRATION

For use in drinking water.

The recommended dosage rates are:

Pigs: 3.33 mg lincomycin and 6.67 mg spectinomycin/kg bw/day, for 7 days. This amounts to 15 mg powder/kg bw/day for 7 days.

Chickens: 16.65 mg lincomycin and 33.35 mg spectinomycin/kg bw/day, for 7 days. This amounts to 75 mg powder/kg bw/day for 7 days.

Pigs:

To determine the volume of dilution (in litres of drinking water) required for 150 g of the veterinary medicinal product, use the following formula:

$$\frac{\text{Volume (L) for 150 g of the veterinary medicinal product}}{10,000 \times \left[\frac{\text{daily water consumption per animal (L)}}{\text{average body weight of one pig (kg)}} \right]}$$

In pigs 150 g of the veterinary medicinal product corresponds to the dose for 10,000 kg of body weight per day.

As an indication, standard water intake varies around 0.15 L/kg bw/day. The table below shows the volume of water to be used for dilution of 150 g of the veterinary medicinal product.

Water consumption	150 g of powder = 100 g antibiotic activity should be diluted in...
0.1 L/kg bw/day	1,000 L of drinking water
0.15 L/kg bw/day	1,500 L of drinking water
0.2 L/kg bw/day	2,000 L of drinking water
0.25 L/kg bw/day	2,500 L of drinking water

Chickens:

To determine the volume of dilution (in litres of drinking water) for 150 g of the veterinary medicinal product, use the following formula:

$$\text{Volume (L) for 150 g of the veterinary medicinal product} = \frac{2,000 \times (\text{daily water consumption per bird (L)})}{\text{average body weight of one bird (kg)}}$$

150 g of the veterinary medicinal product corresponds to the dose for 2,000 kg of body weight per day.

CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substances or any of the excipients.

Do not use in case of hepatic dysfunction.

Do not allow rabbits, rodents (e.g. chinchillas, hamsters, guinea pigs), horses or ruminants to access to water or feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Do not use in laying hens.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals

It is sound clinical practice to base treatment on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of target bacteria.

Use of the veterinary medicinal product deviating from the instructions may increase the risk of development and selection of resistant bacteria and decrease the effectiveness of treatment with macrolides due to the potential for cross-resistance.

The oral use of preparations containing lincomycin is only indicated in swine and chickens.

Do not leave access to the medicated water for other animals. Lincomycin may lead to severe gastrointestinal disturbances in other animal species.

The repeated or prolonged use should be avoided, by improving the farm management and disinfection practices.

Diagnosis should be reconsidered if improvement is not seen after 5 days.

Sick animals have a reduced appetite and an altered drinking pattern, and severely affected animals may therefore require parenteral treatment.

This powder is for use in drinking water only and should be dissolved before use.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to lincomycin, spectinomycin or soybean millfeed should avoid contact with the veterinary medicinal product. Care should be taken not to raise and inhale any dust.

Contact with skin and eyes should be avoided.

Personal protective equipment consisting of approved dust masks, gloves and safety glasses should be worn when handling and mixing the product.

Wash hands and any exposed skin with soap and water immediately after use.

If symptoms such as skin rash or persistent eye irritation appear after exposure, seek medical advice immediately and show the package leaflet or label to the physician.

INTERACTIONS WITH OTHER MEDICATIONS

In general mixture with other medicines should be avoided.

The combination of lincosamides and macrolides is antagonistic, due to competitive binding to their target sites. Combination with anaesthetics may lead to possible neuromuscular blocking.

Do not administer with kaolin or pectine as they impair lincomycin absorption. If co-administration is mandatory, respect a delay of two hours between intakes.

PREGNANCY AND LACTATION

Pigs

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Laboratory studies in dogs and rats have not produced any evidence of reproductive, foetotoxic or teratogenic effects for lincomycin or spectinomycin.

Lincomycin is excreted in milk.

Use only accordingly to benefit-risk assessment by the responsible veterinarian.

Chickens

Do not use in birds in lay.

ADVERSE EFFECTS

Cases of diarrhoea or soft faeces and/or perianal region inflammation have been encountered in healthy pigs at the start of treatment. The symptoms disappeared within 5 to 8 days without interruption of the treatment.

Rare cases of irritability/excitation, skin rash/pruritus were also observed.

Allergic/hypersensitive reactions are rare but can occur and require stopping treatment with the veterinary medicinal product. A symptomatic treatment must be implemented.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

OVERDOSE & TREATMENT

In the event of overdose in pigs, a change in the consistency of the faeces (soft faeces and/or diarrhoea) may be observed.

In chickens treated at several times the recommended dose, enlargement of the caecum and abnormal caecum content was observed.

In case of accidental overdose, the treatment should be interrupted and restarted at the recommended dose.

WITHDRAWAL PERIOD

Pigs:

Meat and offal: Zero days.

Chickens:

Meat and offal: 5 days.

Not authorised for use in birds producing eggs for human consumption, including replacement chicks which are intended to produce eggs for human consumption.

Animals must not be slaughtered for human consumption during treatment.

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in food µg/kg
Lincomycin	Muscle (chicken and pig)	100
	Skin & Fat (chicken and pig)	50
	Liver (chicken and pig)	500
	Kidney (chicken and pig)	1500
	Eggs (chicken)	50
Spectinomycin	Muscle (chicken and pig)	300
	Fat (chicken and pig)	500
	Liver (chicken and pig)	1000
	Kidney (chicken and pig)	5000

PACKING

1kg

LIM SENG PHARMACHEM SDN. BHD.

(product registration holder)

2692, Tingkat Perusahaan Enam,
Prai Industrial Estate Mk 1,

13600 Prai, Pulau Pinang.

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised September 2020