

VP TYLOSIN 50% W/W WATER SOLUBLE POWDER

STRENGTH OF ACTIVE SUBSTANCE

COMPOSITION: per 1g

Tylosin Tartrate..... 547.2mg
(equivalent to Tylosin..500mg)

DESCRIPTION:

Appearance: White to almost white free flowing powder; odour less.

DOSAGE FORM: Powder to be dissolved in water

PHARMACODYNAMICS:

Pharmacodynamic properties Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible microorganisms. The tylosin spectrum of activity includes Gram-positive bacteria, some Gram – negative strains such as *Pasteurella*, and *Mycoplasma spp*

PHARMACOKINETICS:

In most species peak plasma concentrations have been attained 1 to 2 hours after administration of tylosin. Compared to plasma levels clearly higher tissue concentrations have been observed. Tylosin was extensively metabolised. Most of the residues are excreted in faeces predominantly consisting of tylosin A, tylosin factor D and dihydrodesmycosin

TARGET SPECIES:

Bovine (Calves), Poultry (Chicken, Turkeys), Swine(Pigs)

INDICATIONS:

The presence of the clinical disease in the herd/flock should be established before preventive treatment is started.

Pigs:

Treatment and prevention of Porcine Intestinal *Adenomatosis (Ileitis)* associated with *Lawsonia intracellularis*.

Treatment and prevention of enzootic pneumonia caused by *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis*.

Calves: treatment and prevention of pneumonia caused by *Mycoplasma spp.*

Chickens (Broilers – pullets):

Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*

Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens*.

Turkeys: treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum*.

RECOMMENDED DOSAGE :

Oral administration through the drinking water, milk or milk-replacer.

Calves:

10 – 20 mg tylosin per kg BW (corresponding to 20 –40 mg of the VP TYLOSIN 50% per kg BW), twice daily
(= daily dose of 20 – 40 mg tylosin per kg BW), for 7 to 14 days.

Turkeys:

75 – 100 mg tylosin per kg BW per day (corresponding to 150-200 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

Chickens (Broilers, pullets):

For the treatment of chronic respiratory disease:

75 – 100 mg tylosin per kg BW per day (corresponding to 150-200 mg of VP TYLOSIN 50% per kg BW) for 3 – 5 days.

For the treatment of necrotic enteritis:

20 mg tylosin per kg BW per day (corresponding to 40 mg of VP TYLOSIN 50%) for 3 days.

Pigs:

For the treatment of enzootic pneumonia:

20 mg tylosin per kg BW per day (corresponding to 40 mg of VP TYLOSIN 50% per kg BW) for 10 days.

For the treatment of ileitis or PIA:

5 – 10 mg tylosin per kg BW per day (corresponding to 10-20 mg of VP TYLOSIN 50% per kg BW) for 7 days.

For the preparation of the medicated water/milk/milk-replacer the body weight of the animals to be treated and their actual daily water/milk/milk-replacer consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system.

To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{..... mg tylosin / kg bodyweight / day} \times \text{Average body weight (kg) of the animals to be treated}}{\text{Average amount of drinking water / animal (l)}} = \text{.....mg tylosin / l of drinking water}$$

Sufficient access to the system of water supply should be available for the animals to be treated to ensure adequate water consumption.

No other source of drinking water should be available during the medication period.

Should there be no clear response to treatment within 3 days the treatment approach should be reconsidered. After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance which might support development of resistance.

Medicated water, milk or milk replacer should be replaced every 24 hours.

If individual animals show signs of a serious infection such as a reduced water or feed intake, then they should be treated individually, such as by injection.

ROUTE OF ADMINISTRATION:

Oral

CONTRA-INDICATIONS:

Do not use in animals with known hypersensitivity to tylosin or other macrolides.

Do not use in cases with known resistance to tylosin or cross-resistance to other macrolides (MLS-resistance).

Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within 1 week previously.

Do not use in animals with hepatic disorders.

WARNINGS AND PRECAUTIONS

Special warnings (for each target species)

Due to likely variability (time, geographical) in susceptibility of bacteria to Tylosin, bacteriological sampling and susceptibility testing are recommended.

Under-dosing and/or treating for an insufficient length of time are considered to promote the development of resistance in bacteria and should be avoided.

Special precautions for use**Special precautions for use in animals**

Animals with acute infections may have a reduced water and feed consumption and should be treated with a suitable injectable veterinary medicinal product first. The sensitivity of bacteria to tylosin may have changed over time or geographically. It is sound clinical practice to base treatment on susceptibility testing.

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or wildlife.

If individual animals show signs of a serious infection such as a reduced water or feed intake, then they should be treated individually, such as by injection.

Special precautions for the person administering the veterinary medicinal product to animals

Because of the possibility of contact dermatitis and irritation of the skin, eyes or respiratory tract, direct contact during administration should be avoided.

Macrolides may induce hypersensitivity reactions (allergy) after injection, inhalation, ingestion or contact with the skin. Cross-hypersensitivity to macrolides may be observed. Allergic reactions to these substances may be particularly hazardous. Therefore, direct contact during administering of the product should be avoided.

Hypersensitive persons should avoid all contact with the product.

Wear a mask, safety glasses and protective gloves when either reconstituting or administering the solution. After preparation of medicated water, wash exposed skin with soap and water.

In case of accidental eye contact, wash the eyes thoroughly with water.

Contact a physician if a skin rash is observed, in the event of oedema of the face, lips or eyes, or if breathing difficulties are encountered.

INTERACTIONS WITH OTHER MEDICAMENTS:

Antagonism with substances of the lincosamide group

PREGNANCY AND LACTATION:

Laboratory studies in mice and rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

SIDE EFFECTS:

In pigs, adverse reactions have been observed, including diarrhea, pruritus, erythema of the skin, swelling of the vulva, rectal edema and prolapse. These reversible signs appeared 48-72 hours after start of treatment.

SYMPTOMS AND TREATMENTS OVERDOSE:

There is no evidence of tylosin toxicity in rats, at dose rates of up to 1000 mg/kg by the oral route.

There is no evidence of tylosin toxicity in chickens, turkeys, pigs or calves when administered orally at up to three times the recommended dose.

WITHDRAWAL PERIOD:

Meat and offal Pigs:

1 day

Calves: 12 days

Chickens (broilers, pullets) and turkeys: 2 days

Not permitted for use in laying birds producing eggs for human consumption

STORAGE CONDITIONS:

Store below 30°C

Protect from light and moisture

Proposed Shelf Life (as packaged for sale)

2 years as packaged for sale

Proposed Shelf Life (after first opening of container)

NIL

Proposed Shelf life (after reconstitution or dilution)

Any unused solution is discarded after 24 hours

PACKING:

1g x 500 Sachets (Quantity)

MANUFACTURED BY:

Hangzhou An-I-Mate Animal Pharmaceutical Co.,Ltd

No.2 Fengdu Industrial Garden,, , Pingyao,

Yuhang Hangzhou, , , CHINA

NAME & ADDRESS OF PRODUCT REGISTRATION HOLDER**Yeam Heng Trading Co.**

Lot 23, Jalan TPP 5/13, Seksyen 5

Taman Perindustrian Puchong

47100 Puchong, Selangor,

Malaysia.

DATE OF REVISION OF PACKAGE INSERT:

8th July 2020