

PF SALINOMYCIN PREMIX 12% W/W GRANULE

Composition:

Salinomycin Sodium.....120mg

Exp. Qs.....1g

Description

Brown to yellowish-brown granule

Pharmacodynamic

Ionophores facilitate transport of Na⁺ ion in cells and elevates the intracellular concentration of Na⁺ ion. This increased concentration of Na⁺ ion inhibits the certain mitochondrial functions such as substrate oxidation and ATP hydrolysis. Intracellular Na⁺ ion exchanges for extracellular Ca⁺⁺ and increases intracellular concentration of calcium ions lead to cytotoxicity.

Pharmacokinetic

Coccidiostats had a low bioavailability in poultry. Tissue/plasma partition coefficients showed a higher affinity of salinomycin for fat, followed by liver and muscle tissue. The depletion data showed a fairly rapid elimination of coccidiostats in all the tissues after cessation of treatment.

Indication

PF Salinomycin Premix 12% W/W Granule is indicated as an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella* and *E. brunetti* in broiler chickens.

Route of Administration

Oral

Recommended dose

MIXING DIRECTIONS

Thoroughly mix 500 g of PF SALINOMYCIN PREMIX 12% W/W GRANULE with 1000 kg of complete feed to obtain a medicated feed with a salinomycin sodium concentration of 0.0060% (60 mg/kg). The medicated feed can be prepared in the form of meal or pellets.

FEEDING DIRECTIONS

The medicated feed is to be fed continuously as the sole ration up to marketing.

Contraindication

Do not allow turkeys, dogs or horses access to this medicated feed as it is known to be toxic to these species. Extra care should be taken to avoid contamination of feeds for these animals.

Do not use in feeds containing pellet binding agents.

Do not use this feed for treatment of outbreaks of coccidiosis.

Interaction with other medicinal products

Do not administer tiamulin to birds receiving this medicated feed.

Pregnancy and lactation

Do not feed to laying hens producing eggs for Human

Side effects

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

Symptoms and treatment of overdose

Administration of salinomycin sodium in feed at 2 times the recommended dose for 28 consecutive days is safe and nontoxic

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

WARNING

1. Do not feed to replacement, laying or breeding chickens.

2. Operators handling the concentrated product should wear a face mask, gloves and protective clothing.

CAUTION

1. A weight gain reduction may occur in cockerels fed this medicated feed in the absence of coccidial exposure.

2. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two day period.

-Jauhkan ubat dari kanak-kanak/Keep out of reach of children.

Shelf life

24 months

Shelf life after opening

24 hours

Shelf life after reconstitution or dilution

24 hours

Withdrawal period

Chicken (broilers): zero days before slaughter.

Packaging

25kg

Revised date

2/11/2020

Distributed by:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT
MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,
14100 SIMPANG AMPAT, PULAU PINANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

Shandong Qilu King-Phar Pharmaceutical Co., Ltd.

21 Qinglong Road, Pingyin, Jinan, China
Tel: 86-531 83127771 Fax : 86-531 83127774