

# HEMATICEN 200 mg/ml solution for injection

## STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS

Each ml contains:

**Active substances:**

Iron(III)-Ions .....200.0 mg  
as Gletoferron 532.6 mg

**Excipients:**

Phenol .....5.0 mg (as preservative)  
Water for injection, q.s.

Dark brown, slightly viscous solution

## INDICATIONS

For the prevention and treatment of iron deficiency anaemia in piglets.

## CONTRAINDICATIONS

Do not administer to piglets suspected to suffer from deficiency of vitamin E and/or selenium. Do not use in case of hypersensitivity to the active substance or to any of the excipients. Do not use in clinically diseased animals, especially not in case of diarrhoea.

## SIDE EFFECTS

Uncommonly discolouration of the tissue and/or slight, soft swelling may be observed at the site of injection. This should disappear within a few days. Also hypersensitivity reactions can occur. Rarely deaths have occurred in piglets following the administration of parenteral iron dextran preparations. These deaths have been associated with genetic factors or deficiency of vitamin E and/or selenium. Very rarely piglets deaths have been reported which have been attributed to an increased susceptibility to infection due to temporary blocking of the reticuloendothelial system.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s) during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated )
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

## TARGET SPECIES

Pig (piglet)

## DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

For strictly intramuscular injection.

*Piglets:* 200 mg Fe3+ per animal which is equivalent to 1 ml of the product per animal.

Inject once between the 1st and the 3rd day of life.

## ADVICE ON CORRECT ADMINISTRATION

The use of a multidose syringe is recommended. To refill the syringe use a draw-off needle to avoid excessive broaching of the stopper. The stopper must not be broached more than 10 times. When treating groups of animals in one run, use a draw-off needle that has been placed in the vial stopper to avoid excess broaching of the stopper. The draw-off needle has to be removed after treatment.

## WITHDRAWAL PERIOD

Meat and offal: Zero days

## SPECIAL STORAGE PRECAUTIONS

Do not freeze. Do not store above 30°C.

## SPECIAL WARNINGS

Special precautions for use in animals: Not applicable.

Special warnings for each target species: None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to iron dextran, or those with haemochromatosis should avoid contact with the product. Care should be taken to avoid accidental self-injection, as well as contact with the eyes and mouth. In case of accidental injection, seek medical advice immediately and show the package leaflet or label to the physician. Wash hands after use.

Use during pregnancy, lactation or lay:

Not applicable.

Interaction with other medicinal products and other forms of interaction:

The absorption of concomitantly administered oral iron may be reduced. In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal product.

Overdose (symptoms, emergency procedures, antidotes), if necessary:

Transferrin-iron saturation levels leading to increased susceptibility for (systemic) bacterial disease, pain, inflammation reactions as well as abscess formation at the injection site may occur. Persistent discolouration of muscle tissue at the injection site may occur. Iatrogenic poisoning with following symptoms: pale mucous membranes, hemorrhagic gastroenteritis, vomiting, tachycardia, hypotension, dyspnoea, edema of the limbs, lameness, shock, death, liver damage. Supportive measures such as chelating agents can be used.

## SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## OTHER INFORMATION

### Pharmacodynamic properties

Iron is an essential micronutrient. It takes a major role in the oxygen transport of haemoglobin and myoglobin, as well as it has a key role in enzymes, such as cytochromes, catalases, and peroxidases.

Iron has a high recovery rate from metabolism and food ingested. Thus, deficiency occurs only very rarely in adult animals.

### Pharmacokinetic particulars

After intramuscular injection, the iron complex is absorbed into the lymphatic tissue within 3 days. Here, the complex is split to release Fe<sup>3+</sup> which is stored as ferritin in the main storage organs (e.g. liver, spleen and the reticuloendothelial system). In the blood, free Fe<sup>3+</sup> binds to transferrin (transport form) and is mainly used for the synthesis of haemoglobin.

Pack sizes:

- Box with 1 vial of 100 ml
- Box with 1 vial of 200 ml
- Box with 10 vials of 100 ml
- Box with 10 vials of 250 ml

Not all pack sizes may be marketed.

**For animal treatment only. To be supplied only on veterinary prescription.**

**Keep out of the sight and reach of children.**

***"Jauhi ubat dari kanak-kanak"***

***Ubat Terkawal***

**DATE OF REVISION OF LEAFLET**

October 2020

**MARKETING AUTHORISATION HOLDER:**

**Yenher Agro-Products Sdn. Bhd.**

1628, Jalan IKS Simpang Ampat 1, Taman IKS  
Simpang Ampat, 14100 Simpang Ampat, Seberang  
Perai Selatan, Penang, Malaysia.

**MANUFACTURER:**

**CENAVISA S.L.**

Camí Pedrá Estela, s/nº, Reus,  
43205, Tarragona España

 **cenavisa**  
**LABORATORIOS**

Tel 34-977 75 72 73

[www.cenavisa.com](http://www.cenavisa.com)

E-mail: [cenavisa@cenavisa.com](mailto:cenavisa@cenavisa.com)