

DOTFEED 250G/KG POWDER

Description and Composition

DOTFEED 250G/KG POWDER is an off-white to pale cream powder. Each kg contains 250g of Dinitolmide as its active ingredient.

Pharmacodynamics

Dinitolmide is a nitrobenzamide anticoccidial agent. It acts primarily on the first-generation schizonts, and it inhibits sporulation of oocysts. It is coccidiostatic if given for 6 days but is coccidiocidal if given for longer periods.

Pharmacokinetics

Benzamide anticoccidials include dinitolmide (also called zoalene), aklomide and nitromide. Of these, dinitolmide is most widely used for treating coccidiosis in poultry. Dinitolmide is rapidly absorbed following oral administration to chickens, and broadly and rapidly distributed to tissues. Following absorption, dinitolmide is extensively metabolized by the liver and excreted in faeces.

Indication

An aid in the prevention of coccidiosis of broilers, replacement pullets and turkeys.

Recommended Dose

Broilers and replacement pullets: Thoroughly mix 500g of product per tonne of feed to produce a concentration of 125g Dinitolmide per tonne in the finished feed.

Turkey growers: Thoroughly mix 760g of product per tonne of feed to produce a concentration of 190g Dinitolmide per tonne in the finished feed. Use continuously for up to 14 days, or as directed by a veterinarian.

Route of Administration

For oral administration via feed.

Contraindication

Birds receiving this medication should not be treated with Nitrofurans (eg. Nitrofurazone, Furaltadone).

Warning and Precautions

Special warning for each target species:

Nil

Special precautions for use in animal:

Do not use in laying hens (chickens/turkeys) or in replacement pullets over 14 weeks of age.

Interaction with Other Medicaments

Birds receiving this medication should not be treated with Nitrofurans (eg. Nitrofurazone, Furaltadone).

Pregnancy and Lactation

The single study available on the deposition of dinitolmide in eggs did not continue sampling eggs long enough to establish the full time course of drug elimination, but drug residues were still present 2 weeks after the medication was withdrawn (Nose *et al.*, 1982). Concentrations of dinitolmide residues were approximately 10 times higher in the egg yolk than the albumen (Nose *et al.*, 1982).

Side Effects

No adverse effects are seen when the drug is used as directed.

Symptoms and Treatment of Overdose

The severity of the toxic effects of zoalene on baby chicks is dose dependent. In laying birds however, there was cessation of egg production, capricious appetite and progressive weakness, but no nervous signs or deaths, even when as much as 10 times the recommended level was fed for 30 days. Mortality was not a feature of zoalene poisoning when feeding as much as 10 times the recommended level to baby chicks.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Period

Meat: Nil

Eggs: Do not use in laying hens (chickens/turkeys) or in replacement pullets over 14 weeks of age.

Packing

10kg and 25kg.

Revision date: 3.9.20

	Manufacturer & Product Registration Holder Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com
---	--