

Allercrom Eye Drops 2% w/v

(Sodium Cromoglycate 2% w/v)

Description

Clear colorless solution free from visible particulate matter.

Composition

Each ml contains 20 mg of cromoglycic acid (INN) in the form of sodium salt (sodium cromoglycate).

Excipients: s.q. (0.01% benzalkonium chloride as preservative).

Pharmacodynamics

Cromoglycic acid (sodium cromoglycate) inhibits the degranulation of sensitized mast cells, preventing the release of histamine and other inflammation mediators. It has none of the collateral of the antihistamines, corticosteroids, anticholinergic or vasoconstricting agents. It is practically innocuous and can be combined with other preparations administered by topical or systemic route.

Pharmacokinetics

Sodium cromoglycate is poorly absorbed. When multiple doses of Sodium cromoglycate ophthalmic solution are instilled into normal rabbit eyes, less than 0.07% of the administered dose of Sodium cromoglycate is absorbed into the systemic circulation (presumably by way of the eye, nasal passages, buccal cavity and gastrointestinal tract). Trace amounts (less than 0.01%) of the Sodium cromoglycate does penetrate into the aqueous humour and clearance from this chamber is virtually complete within 24 hours after treatment is stopped.

In normal volunteers, analysis of drug excretion indicates that approximately 0.03% of Sodium cromoglycate is absorbed following administration to the eye.

Indications

Prevention and relief of the ocular symptoms that appear with allergic conjunctivitis and seasonal conjunctivitis.

Recommended Dose

Generally instill 1 drop of eye drops into each eye 4 times daily, at regular intervals. If symptoms worsen with the use of the product, discontinue the use of the medication and consult physician.

Route of Administration

Separate the eyelids from the eye and instill the drops in the conjunctival sac.

Application of the eye drops should be carried out under hygienic conditions, avoiding contact of the dropper tip to any surface.

Close bottle after every application.

Discard one month after first opening the bottle.

Contraindications

Hypersensitivity to the components of the formula.

Warning and Precautions

This drug should be used during pregnancy only if clearly needed. Caution should be exercised when it is administered to a nursing woman.

Interactions with other medicaments

It has not been described that Allercrom Eye Drops 2% w/v presents interaction with any other drug.

Pregnancy and Lactation***Pregnancy***

As with all medication, caution should be exercised especially during the first trimester of pregnancy. Cumulative experience with Sodium cromoglicate suggests that it has no adverse effects on foetal development. It should be used in pregnancy only where there is a clear need.

Lactation

It is not known whether Sodium cromoglicate is excreted in human breast milk but, on the basis of its physicochemical properties, this is considered unlikely. There is no information to suggest the use of Sodium cromoglicate has any undesirable effects on the baby.

Side Effects

It may cause stinging or irritation that disappears during the first days of treatment.

Symptoms and Treatment of Overdose

Due to the characteristics of this preparation, intended for topical use, no toxic effects are expected with the use of this product at the recommended doses or in the event of accidental ingestion of the contents of a bottle.

Storage Condition

Protect from light. Store below 30°C.

Shelf-life

24 months from date of manufacture for unopened vial and 4 weeks from the date of opening the vial.

Presentation

5 ml solution fill in 5 ml, 10 ml solution fill in 10 ml and 13.5 ml solution filled in 15 ml labeled LDPE vial with spike HIPS cap packed in a carton.

Revised date: 1/6/2017

Manufactured by:

FDC Limited
B-8, MIDC Area,
Waluj, Aurangabad,
431 136 India.