

Important information. Please read carefully.

AsmafenTM

Ketotifen 1mg

COMPOSITION

Asmafen Tablet: Each tablet contains Ketotifen (as hydrogen fumarate) 1mg

INDICATIONS

Asmafen is indicated for long-term prevention of bronchial asthma, allergic bronchitis, asthmatic symptoms associated with hay fever. It is also indicated for the prevention and treatment of other allergic conditions such as allergic rhinitis, conjunctivitis and allergic skin disorders.

PHARMACODYNAMICS

Ketotifen Fumarate has the properties of the antihistamines in addition to actions on mast cells. Its mode of action is not completely understood but it prevents release of pharmacological mediators of bronchospasm by stabilising mast cell membranes.

PHARMACOKINETICS

Ketotifen Fumarate is absorbed from the gastrointestinal tract and together with its metabolites, is excreted in the urine and faeces.

DOSAGE AND ADMINISTRATION

To be administered orally. Unless otherwise prescribed by the physician:

Adults and children above 12 years	:	1 to 2mg (twice a day)
Children 2 to 12 years	:	1mg (twice a day)

CONTRAINDICATIONS

Asmafen is contraindicated for use during acute attacks of asthma and in patients who are on oral antidiabetic treatment.

PRECAUTIONS

Do not discontinue treatment abruptly as this may lead to recurrence of the symptoms of asthma. Should withdrawal be necessary, reduce the dose gradually over a period of one week. Avoid alcoholic drinks.

PREGNANCY AND LACTATION

Use in pregnancy is best avoided. When necessary, close medical supervision is required.

DRUG INTERACTIONS

Ketotifen may potentiate the sedative effects of other CNS depressants, antihistamines and alcohol.

SIDE EFFECTS

This preparation may cause drowsiness. Patients should be advised against driving motor vehicles or operating machinery. Other side effects such as dry mouth, dizziness and weight gain may also occur.

OVERDOSAGE AND TREATMENT

Overdosage may lead to drowsiness, confusion, dyspnoea, bradycardia or tachycardia, disorientation, and convulsions. If overdosage is recent, the stomach should be emptied by gastric lavage. Other supportive treatments should also be given if necessary.

PRESENTATION

White, round, flat bevelled edge tablet with one breakline and 'XS' embossed on one side, 7.0 mm in diameter in blister packs of 100's.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE

Protect from light. Store below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 6-Nov-2013

Manufactured and Marketed by
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

AS0503-001

11.13

