

Florfenicol 25%w/v
Oral Solution
(FOR VETERINARY USE ONLY)

COMPOSITION

Florfenicol 250 mg
Exp q.s to 1 ml

PRODUCT DESCRIPTION

The finished product is a clear yellowish solution.

TARGET SPECIES

Pigs

PHARMACODYNAMIC

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

PHARMACOKINETIC

After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

INDICATIONS

Treatment and prevention at the group level where clinical signs are present of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol. The presence of the disease should be established in the herd before initiating preventive treatment.

WARNINGS AND PRECAUTIONS

1. Special warnings for each target species:

The treated pigs should be placed under special observation. On each of the five days of treatment, unmedicated drinking water should not be given until the full daily amount of medicated drinking water has been ingested by pigs. If there are no signs of improvement after three days of treatment, the diagnosis should be reviewed and, if necessary, the treatment changed.

2. Special precautions for use in animals:

The veterinary medicinal product should be used in conjunction with susceptibility testing.

Use of the product deviating from the instructions given may increase the prevalence of bacteria resistant to the florfenicol. Official and local antimicrobial policies should be taken into account when the product is used.
Treatment should not exceed 5 days.

3. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to florfenicol or polyethylene glycols should avoid contact with the veterinary medicinal product. In case of accidental spillage onto skin rinse with water. In case of contact with eyes, rinse immediately with copious amounts of water. Personal protective goggles should be worn when handling the veterinary medicinal product.

4. Other precautions:

In order to prevent any adverse effects on algae and possible contamination of groundwater, manure from treated pigs must not be spread onto land without dilution with manure from untreated pigs. Manure from treated pigs must be diluted with at least 5 times the weight of manure from untreated pigs before it can be spread onto arable land.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

CONTRA-INDICATIONS

Do not use in boars intended for breeding purposes.

Do not use in case of hypersensitivity to the active substance or to the excipient.

ADMINISTRATION & DOSAGE

To be administered in drinking water.

10 mg florfenicol per kg bodyweight per day in drinking water for 5 consecutive days. This dose is equivalent to 0.04 ml of the product per kg body weight per day.

Based on the recommended dose, and the number and weight of the animals to be treated, the exact daily amount of the veterinary product should be calculated according to the following formula:

X ml veterinary product/kg b.w./day	x	Mean body weight (kg) of animals to be treated	=	X ml veterinary product per liter drinking water
Mean daily water consumption (litre) per animal				

The appropriate quantity of medicated water should be prepared based on the daily water consumption. To ensure a correct dosage body weight should be determined as accurately as possible. In order to avoid under- and over-dosing, treated animals should be divided into groups of similar bodyweight and the dose should be calculated for each group individually.

INTERACTION WITH OTHER MEDICAMENTS

No data available

PREGNANCY AND LACTATION

Studies in laboratory animals have not revealed any evidence of potential embryotoxic or foetotoxic effect of florfenicol. The safety of the veterinary medicinal product in sows has not been established during pregnancy and lactation.
The use is not recommended during pregnancy and lactation.

OVERDOSE AND TREATMENT

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

SIDE EFFECT

A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment. Commonly observed adverse effects are diarrhoea and/or peri-anal and rectal erythema/oedema which may affect approximately 40% of the animals. These effects are transient. In a few of the affected animals, prolapse of the rectum, that resolves without treatment may be observed.

WITHDRAWAL TIME

Meat and offal: 20 days

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store in a cool dry place ($\leq 25^{\circ}\text{C}$). Strictly avoid light & heat exposure.

PACKAGING

The product filled 1000 ml in HDPE plastic bottle, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 07 Jun 2019



Product Registration Holder and Manufactured By:
SHENNONG ANIMAL HEALTH (M) SDN BHD
神農動物保健(馬)有限公司
No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,
Taman Perindustrian USJ 1,
47610 Subang Jaya, Selangor, Malaysia.