

AMPROLIUM 20% W/W WATER SOLUBLE POWDER

Amprolium 20% w/w Water Soluble Powder appears to be a light yellow color powder.

Composition

Each gram contains Amprolium (as Hydrochloride) 200 mg

Pharmacodynamics

Amprolium is an anticoccidial agent that acts as a competitive inhibitor of thiamine in the parasite metabolism, and interferes with the metabolism of glucides necessities for coccidian multiplication and survival.

In in-vitro studies, it was shown that the uptake of thiamine by schizonts of *Eimeria tenella* and by host intestinal cells can occur through passive diffusion or by an active, energy- and pH-dependent process. Amprolium competitively inhibited both systems, however, the parasite was shown to be more sensitive to amprolium than the host.

As shown with *Eimeria maxima* inoculated chicken, the administration of Amprolium resulted in a proportion of morphologically abnormal macrogametes and oocysts which may be considered the reason for a reduced sporulation rate.

Pharmacokinetics

Absorption is low after oral administration as maximum concentration was achieved only after 4 hours. Excretion is mainly through faeces.

Indication

For the treatment of coccidiosis in growing chickens, turkeys and laying hens.

Dosage and administration

Oral administration.

Administer 100g of product to 170.55 liters of drinking water as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, administer 200g of product to 170.55 liters of drinking water). Continue with 50g of product to 170.55 liters for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as sole source of amprolium.

Contraindication

None known.

Warning and Precautions

Special warnings for each target species

As with any antiparasiticide, frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance. In case of detection on the lack of efficacy during treatment, do communicate to the national competent authorities.

Special precautions for use

(i) Special precautions for use in animals

The product is not intended for a preventive use.

This product should be reserved in case of coccidiosis outbreaks due to non-availability of vaccine, in case of lack of efficacy of vaccine and in vaccinated flocks if a severe coccidial challenge is diagnosed before immunity has fully developed.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

This is an irritant and corrosive product. It could cause airway, eye and skin irritation.

Wear impervious gloves and protective glasses when handling the product.

Avoid inhalation of vapours.

Avoid contact with skin and eyes. If skin contact occurs remove contaminated clothing and wash skin thoroughly with soap and water.

Seek medical attention if irritation persists.

This product is harmful when ingested. In case of accidental ingestion, rinse mouth with water. Give plenty of water to drink. Do not induce vomiting and seek medical attention immediately.

Person with known hypersensitivity to the active substance or to any of excipients should avoid contact with the product.

Wash hands and exposed skin after use.

Interactions with Other Medicaments

Amprolium is a thiamine analogue. Therefore, the efficacy of amprolium may be reduced during a simultaneous administration of products containing vitamin B-complex.

Statement on Usage during Pregnancy and Lactation

Studies in animal laboratory have not produced any evidence of teratogenic effects. The safety of amprolium has not been investigated in laying birds. Use only according to the risk/benefit assessment by the responsible veterinarian.

Side Effects

None known.

Overdose and Treatment

Prolonged use can produce thiamine deficiency. If the symptoms appear, thiamine should be administered.

Withdrawal Period (s)

Nil.

Storage Conditions

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/Jauhkan daripada kanak-kanak.

Any unused product or packaging should be disposed of in accordance with local legislation.

Shelf-life after opening: 7 days.

Shelf-life after reconstitution: 24 hours.

Packaging Available

100 G & 1 KG.

Product Holder/ Manufacturer

NAM PHARMA SDN BHD

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Date of revision: 30-05-2024