



RHOMICOSIN® 20FG Oral Powder 20%w/w

Composition: Tilmicosin 20%w/w (equivalent to 22.25% w/w of tilmicosin phosphate)

Product Description: White to off-white powder

Pharmacodynamics: Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal. This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular its activity has been demonstrated against the following micro-organism:

Pigs: *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*. Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit *in vitro* the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion. Cross resistance between tilmicosin and other macrolides and lincomycin has been observed.

Pharmacokinetics:

Absorption: When administered to pigs via the oral route at a dose of 400 mg tilmicosin/kg feed (equivalent to approximately 21.3 mg tilmicosin/kg bodyweight/day), tilmicosin moves rapidly out of the serum into areas of low pH.

Distribution: Following oral administration, tilmicosin is distributed throughout the body with especially high levels found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues.

Biotransformation: Several metabolites are formed, the predominant one being identified as T1. However the bulk of tilmicosin is excreted unchanged.

Elimination: Following oral administration, tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine.

Indication:

Pigs: Prevention and treatment of respiratory disease caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin.

Routes of administration: Oral

Dosage: Administer in the feed at a dose of 8 to 16 mg/kg body weight/day of tilmicosin (equivalent to 200 to 400 ppm in the feed) for a period of 15 to 21 days. Inclusion rate is 1-2 kg of product per tonne of feed.

To ensure thorough dispersion of the product, it should first be mixed with a suitable quantity of feed ingredients (10 kg) before incorporation into the finished feed.

Contraindications: Horses or other Equidae, must not be allowed access to feeds containing tilmicosin. Horses fed with tilmicosin medicated feeds may present signs of toxicity with lethargy, anorexia, reduction of feed consumption, loose stools, colic, distension of the abdomen and death. Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Warnings and Precautions:
Special precaution for target species

Under practical conditions, the management of respiratory disease outbreaks recognises that acutely ill animals are inappetent and require parenteral therapy.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator or a non-disposable respirator. Do not eat, drink or smoke when handling this product. Wash hands after use.

In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Special precautions for disposal of unused products

Drug containers and any residual contents should be disposed of safely, according to the regulation in force.

Interactions with other medicaments: None known

Statement on usage during pregnancy and lactation: The safety of tilmicosin has not been established in boars used for breeding purposes.

Adverse Effects: In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

Overdose and Treatment: No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

Storage Conditions: Store in dry place below 30°C. Protect from direct sunlight.

Shelf life after reconstitution: To be used within 24 hours.

Withdrawal Period:

Pigs: 21 days

FOR VETERINARY USE ONLY

Registration holder and manufactured by:



RHONA MA MALAYSIA SDN BHD (200001023279)

Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia.

Tel: 03-7873 7355 Fax: 03-2770 0119

Email: customercareline@rhonema.com

Date of revision: February 2020
5LRHOMI3-10K 02/FEB20

W10cm H14cm

LABEL CODE:	VERSION	COLORS
5LRHOMI3-10K PRODUCT: RHOMICOSIN 20 FG ORAL POWDER 20% W/W PACK SIZE: 10 KG COUNTRY: MY DIMENSIONS: cm = 10 x 14	Version: 2 MMM/YY: FEB 20	Black ! Please use the official PANTONE® (P) matching • system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME: DATE + SIGNATURE:	COMPLETE NAME: DATE + SIGNATURE:	COMPLETE NAME: DATE + SIGNATURE: