



RHOBAZIN™ PLUS

Solution for
Oral Use

Composition :

Sodium sulfaquinoxaline: 5.37%w/v
(equivalent to 5% w/v sulfaquinoxaline)
Pyrimethamine 1.5% w/v

Product Description :

A yellow amber viscous solution, miscible with water.

Pharmacodynamics :

Pyrimethamine is a pyrimidine derivative, folic acid antagonist, and acts as an inhibitor of dihydrofolate reductase. The sulphaquinoxaline is a sulfonamide that acts as an antagonist of PABA, suppressing the synthesis of tetrahydrofolic acid essential to the protozoa. Both compounds are synergistic. Its association is highly effective prophylactically and therapeutically for coccidiosis. The ratio for both active ingredients is 1:3.

They are active against coccidia as follows: Chicken: *Eimeris tenella*, *E. necatrix*, *E. bruneti*, *E. acervulina*, *E. maximum*, *E. mivati*, *E. praecox*, *E. Hagani*, *E. mitis*. Turkey: *E. meleagridis*, *E. adenoids*, *E. gallopavonis*.

Pharmacokinetics :

Pyrimethamine has very slow drug absorption. Its average life is 1.5 days via oral administration and being eliminated by the kidneys.

The sulphaquinoxaline is rapidly absorbed orally through the gastrointestinal tract. The degree of absorption varies with species. It is widely disseminated in the tissues and cerebrospinal fluid. The maximum concentration in blood reaches within 3-5 hours. It is metabolized via acetylation in the liver and is excreted in urine for partly unmetabolized, with an average disposal time of 11 hours.

Indication :

Chicken and Turkey: Treatment of cecal and intestinal coccidiosis.

Dosage and Administration:

Oral, in drinking water.

Chicken and Turkey: 15-22.5mg pyrimethamine + 50-75mg sulfaquinoxaline per liter of drinking water (equivalent to 1-1.5 ml Rhobazin™ Plus/litre of drinking water) for three days, stop for 2-3 days and resume treatment for another three days at the same dose.

Contraindication:

Do not administer to:

- Animals with haemorrhages
- Animals with liver or kidney impairment.
- Adult breeding poultry

Warning & Precaution:

Special precautions for use

Water abundantly during treatment.

Handle this product with care and avoid direct contact.

Take all recommended precautions by wearing protective clothing, glasses, gloves and

mask. Avoid contact with skin and eyes.

In case of symptoms as cutaneous eruption, consult a doctor and show him this information. An oedema at the level of the face, the eyelids, the lips or respiratory difficulties constitute severe symptoms and require urgent medical cares. Avoid to handle this product or the supplemented water in case of allergy to sulphonamides.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interaction with other medicaments :

It is antagonistic with PABA-like compounds, such as local anesthetics procaine type: butacaine, benzocaine and procaine penicillin.

Statement on usage during pregnancy & lactation:

Do not use in adult breeders poultry and layers.

Adverse Effect:

Prolonged treatment (over 5 days) acute hemorrhage may occur in some treated chickens, especially in warm seasons. In this case, supplementation of vitamin K is required.

Overdose and Treatment:

If overdosed, diarrhea and sagging may present. To discontinue administration.

Storage Condition:

Store in dry place below 30°C. Protect from direct sunlight.

Withdrawal Period:

Treated birds must not be slaughtered for use in food for at least 4 days after the latest treatment.

Packing: 1L and 5L.

Registration holder and manufactured by:



RHOMA MALAYSIA SDN BHD (200001023279)

Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia.

Tel: 03-7873 7355 Fax: 03-2770 0119

E-mail: customer@rhonema.com

Date of revision: JANUARY 2020
5LRHOB3-01L 03/JAN20

LABEL CODE:	VERSION	COLORS
5LRHOB3-01L		Black
PRODUCT: RHOBAZIN PLUS SOLUTION FOR ORAL USE	Version: 3	
PACK SIZE: : 1L , 5L	MMM/YY: JAN20	
COUNTRY: MY		
DIMENSIONS: cm = 10 X 14		! Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:	DATE + SIGNATURE: