

Insert PUP

draft -01

07 Sep 17

PULCO

PREMIX POWDER 20%

Ingredient(s):

Each gram contains:

Tilmicosin Phosphate (eq. to Tilmicosin 200mg) 222.55mg

Pharmacodynamics:

Like other macrolides, tilmicosin has activity primarily against gram-positive bacteria, although some gram-negative bacteria are affected and the drug reportedly has some activity against mycoplasma. Preliminary studies have shown that 95% of studied isolates of *Pasteurella haemolytica* are sensitive.

Pharmacokinetics:

Tilmicosin apparently concentrates in lung tissue. At 3 days post injection, the lung serum ratio is about 60:1. MIC₉₅ concentrations (3.12 micrograms/mL) for *P. Haemolytica* persist for a minimum of 3 days after a single injection.

In pigs, after administration of a single oral dose of tilmicosin, more than 90% of the dose was excreted within 6 days, and 96% of the dose was excreted within 13 days (about 80% in feces and 15% in urine).

Indication:

For control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*

Target Species:

Swine

Recommended Dose:

Thoroughly mix 0.91-1.82 kg Pulco Premix Powder 20% into each ton feed (to provide 181.8g – 363.6 g tilmicosin activity per tonne in complete feed).

Feed for a 21-day period, beginning approximately 7 days before an anticipated disease outbreak. Feed continuously as the sole ration.

Mode of Administration:

Oral via feed mixing.

Contraindication:

1. Do not use in animals hypersensitive to tilmicosin.
2. Tilmicosin is known to be toxic for horses.

Warning and Precautions:

1. The safety of tilmicosin has not been established in boars used for breeding.
2. Do not use in any feed (supplement, concentrate, or complete feed) containing bentonite.
Bentonite in feeds may affect the efficacy of tilmicosin.
3. Do not allow horses or other equine access to feeds containing tilmicosin.

Interaction with Other Medicaments:

1. Do not use in any feeds containing bentonite. Bentonite in feeds may affect the efficacy of tilmicosin.
2. In swine, epinephrine increased the mortality associated with tilmicosin.

Pregnancy and Lactation:

Safe use in pregnant animals or animals to be used for breeding purposes has not been demonstrated.

Side Effects:

In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

Environmental Property:

No environmental data for the mixture or formulation. The environmental information for Tilmicosin is presented.

Tilmicosin is practically nontoxic to fish, birds, earthworms, fungus, molds, soil bacteria, and most plants. Slightly toxic to aquatic invertebrates. Moderately toxic to nitrogen-fixing bacteria. Highly toxic to green algae and blue-green algae. No volatility expected. Low potential to bioconcentrate in aquatic organisms. Low mobility in soil. Persistent in the soil environment. Persistence in the aquatic environment not expected due to rapid photolysis.

Insert PUP
draft -01
07 Sep 17

Symptoms and Treatment of Overdose:

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilimicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

Storage Condition:

Store at temperature below 30°C. Protect from direct sunlight and water.

Shelf Life:

3 years.

Withdrawal Periods

7 days.

Maximum Residual Limit (MRL):

Swine

Muscle, Fat: 50ppm.

Liver, Kidney: 1000 ppm.

Product Description:

Beige yellowish powder.

1kg aluminum pouch and 10kg drum

To declare Y.S.P. INDUSTRIES VIETNAM CO., LTD.
as Manufacturer and Y.S.P. INDUSTRIES (M) SDN.
BHD. as Product Registration Holder

**Made by YSP**

Manufacturer:

Y.S.P. INDUSTRIES VIETNAM CO., LTD

Road No.6, Nhon Trach 3 Industrial zone, Nhon Trach,
Dong Nai province, Vietnam Tel: +8461 3569 993

Product Registration Holder :

Y.S.P. INDUSTRIES (M) SDN. BHD.(192593 U)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679

To update the Date of
revision

Date of revision: 07 Sep 17