

FLUMECIN 20% SOLUTION

Description and Composition

Flumecin 20% Solution is a faint yellow clear solution. Each ml contains 200mg of Flumequine

Pharmacodynamics

Flumequine is an anti-infective synthetic quinolone. It acts on the nucleic acid level by blocking the DNA replication of the bacteria. The molecular target is the DNA gyrase, which is involved in the spiral winding of the DNA.

Flumequine has a dose-dependent bactericidal activity against Gram-negative bacteria. It also has activity on certain Gram-positive bacteria (including staphylococci).

Pharmacokinetics

Flumequine is often well absorbed after administration of the product orally. Its bioavailability is about 76-100% in calves, 57% in chicken, and 7% in pigeon. Tmax is reached after about 2-3 hours and Cmax varies by species and specialty.

The rate of plasma protein binding of flumequine is relatively high (75% in calves and 39% in pigeons).

Pharmacokinetics parameters are linearly correlated with the dose of flumequine administered. No accumulation in the blood, or induction or enzyme inhibition is observed.

Flumequine is metabolized to 7-hydroxyflumequine and glucuronide flumequine in the liver. It undergoes a phase of relatively rapid elimination.

Half-life of elimination is about 1.5 hours in the pigeon, 2 hours in calves, and 7 hours in chicken.

Flumequine is excreted in the urine as flumequine and glucuronide flumequine.

Indication

Poultry: For the treatment and prevention of respiratory infections, gastrointestinal infections and septicemia caused by *Escherichia coli*, *Salmonella spp.* and *Pasteurella spp.*

Calves:

For the treatment and prevention of gastrointestinal infections caused by *Escherichia coli* and *Salmonella spp.* and respiratory infections due to *Pasteurella multocida* and *Mannheimia haemolytica*.

Recommended Dose

Poultry and calves:

12mg flumequine per kg body weight per day for 5 days orally.

The amount of medicated drinking water consumed by the animals depends on their physiological and clinical status.

Route of Administration

For oral administration in drinking water.

Contraindication

Do not use in case of hypersensitivity to quinolones.

Do not use in laying birds.

Warning and Precautions

Special warnings for each target species

None.

Special precautions for use in animals

-The use of quinolones should be based on susceptibility testing whenever possible, and take into account official and local policies of antibiotic use.

-The use outside the recommendations of the SPC may increase the prevalence of bacteria resistant to quinolones and may decrease the effectiveness of treatment with other (fluoro) quinolones due to the potential for cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

-People with known hypersensitivity to quinolones should avoid any contact with the product.

-In case of contact with eyes, rinse immediately with plenty of water. If irritation persists, consult a physician.

Other precautions

None.

Interaction with Other Medicinal Product

Do not combine with trimethoprim or sulfonamides.

Pregnancy and Lactation

Not applicable

Side Effects

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

Symptoms and Treatment of Overdose

None known.

Storage Condition

Store below 30°C.

Shelf life

3 years. To be used within 24 hours after dilution. After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Periods

Poultry and calves:

Meat and offal: 2 days

Eggs: 4 weeks

Packing

1Liter

REVISED: 29.3.16

	Manufactured by::
	Range Pharma Sdn Bhd 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@ rangepharma.com.my