

For animal use only

EAGLEVET FLO 4% (W/W) POWDER

[Active Ingredient]

Florfenicol40g/kg

[Product Description] Yellowish powder

[Pharmacodynamics] Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic. However, bactericidal activity has been demonstrated *in-vitro* against *Pasteurella multocida* when florfenicol is present at concentrations above the MIC for 4 to 12 hours.

[Pharmacokinetics] After administration to pigs by gavage at 10 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5µg/ml were reached approximately 3 hours after dosing. The terminal half-life was between 3 and 4 hours. When pigs were given free access, for 5 days, to feed medicated with florfenicol (premix for medicated feeding stuff) at the recommended dose of 10 mg/kg serum florfenicol concentrations exceeded 1µg/ml for more than 16 hours each day of treatment. Florfenicol is well absorbed when administered orally and following distribution it is rapidly eliminated in the urine and faeces in a ratio of 3:1. A fraction is excreted unchanged and the rest is metabolized into 5 major metabolites.

[Indication] In fattening pigs: For the treatment and prevention of swine respiratory disease in infected herds due to *Pasteurella multocida* susceptible to florfenicol. The presence of the disease should be established in the herd before initiating preventive treatment.

[Recommended Dosage and Administration]

Dosage:

10 mg of florfenicol per kg body weight (bw) (equivalent to 250 mg the veterinary medicinal product) per day administered for 5 consecutive days.

Administration:

For a daily feed intake of 50 g/kg bodyweight, this dosage corresponds to a rate of incorporation of 5 kg of medicated premix per ton of feed, i.e. 200 ppm of florfenicol.

The rate of incorporation of the medicated premix in the feed may be increased in order to achieve the required dosage on a mg/kg bodyweight basis and to take into account the actual feed intake.

Thus, the inclusion level may need adjusting as follows to give the correct dose.

[Contraindications]

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in case of known resistance to florfenicol.

[Warnings and Precautions]

i) Special precautions for use in animals

The product should be used in conjunction with susceptibility testing and take into account official and local policy relating to the use of antimicrobials. This premix is intended for the manufacturing of solid medicated feed and cannot be used as is; the incorporation rate of the premix in feed cannot be lower than 5kg/ton. This premix contains ground limestone, which can lead to a decrease in food consumption and to a phosphorus calcium imbalance in feed intake. Care should therefore be taken to consider the calcium content of the final medicated feeding stuff. Treatment should not exceed 5 days.

ii) Special precautions to be taken by the person administering the medicinal product to animals

Skin sensitization may occur. Avoid skin contact. Handle this product with care to avoid exposure during incorporation of premix into feed and administration of medicated feeding stuff to animals, taking all recommended precautions. Wear gloves and do not smoke, eat, or drink when handling the product or medicated feeding stuff. Wash hands thoroughly with soap and water after use of the product or medicated feeding stuff. Rinse thoroughly with water in case of exposure. If you develop symptoms following exposure such as skin rash, you should seek medical advice immediately and show and take the package leaflet or the label to the physician.

[Interactions with Other Medicaments] No data available

[Pregnancy & Lactation] The safety of the veterinary medicinal product has not been established in sows during pregnancy and lactation. Toxicity studies in rats have shown adverse effects on the male reproductive system. Do not use in pregnant and lactating sows. Do not use in breeding boars.

[Adverse Effects/Undesirable Effects] Commonly observed adverse effects are diarrhoea perianal inflammation and rectal eversion. These effects are transient, resolving on cessation of treatment. Increased serum calcium may also be observed.

[Overdose and Treatment] In the event of overdose, a reduction in food and water consumption, together with a decrease in bodyweight may be observed. There may be an increase in refused feed and an increase in serum calcium.

[Withdrawal Period(s)] Meat and offal - 14 days

[Dosage Forms and packaging available] Powder form, 10 kg (10 x 1 kg ea)

[Date of Revision of Package Insert] 16 August 2016

Name and address of manufacturer



EAGLE VET TECH CO. LTD

Company address: 8th F, Eagle Town Bldg, 278-20, Sungsu-2Ka, Sungdong-Ku, Seoul, Korea
Plant address: 235-34, Chusa-Ro, Shinam-Myun, Yesan-Kun, Chungchongnam-Do, Korea

Name and address of marketing authorization holder

QL Pacific Vet Group Sdn Bhd

No 886-C8, Jalan Subang 9, Taman Perindustrian Subang,
47600 Subang Jaya Selangor, Malaysia

KEEP OUT OF REACH OF CHILDREN / JAUHI DARI KANAK-KANAK
FOR ANIMAL USE ONLY / UNTUK KEGUNAAN HAIWAN SAHAJA
CONTROLLED MEDICINE/ UBAT TERKAWAL