



Controlled Medicine
Veterinary Use Only

Kenoxin Enrofloxacin 10% w/v Oral Solution

1. Brand or Product Name / Name and Strength of Active Substance(s)

Kenoxin Enrofloxacin 10% w/v Oral Solution(Each 1L contains)
Enrofloxacin ----- 100g

2. Product Description : Pale yellowish to yellow liquid

3. Pharmacodynamics/Pharmacokinetics/Environmental Properties

Pharmacodynamics :

Enrofloxacin is a bactericidal agent. Its mechanism of action is believed to act by inhibiting bacterial DNA-gyrase, thereby preventing DNA supercoiling and DNA synthesis.
Both enrofloxacin and ciprofloxacin have similar spectrums of activity. These agents have good activity against many gram-negative bacilli and cocci, including most species and strains of Pseudomonas aeruginosa, Klebsiella spp, E.coli, Enterobacter, Campylobacter, Shigella, Aeromonas, Haemophilus and Proteus. Of the currently commercially available quinolones, ciprofloxacin and enrofloxacin have the lowest MIC values for the majority of these pathogens treated.
The fluoroquinolones have variable activity against most streptococci and are not usually recommended for use in these infections. These drugs have weak activity against most anaerobes and are ineffective in treating anaerobic infections.

Pharmacokinetics:

Enrofloxacin administered via drinking water to poultry is rapidly and very well absorbed with a bioavailability of approx. 90%. Maximum plasma concentrations of 2 mg/L are reached within 1.5 hours after a single bolus dose rate of 10 mg/kg body weight with a total systemic availability of 14.4 mg.hr/L.
Enrofloxacin is eliminated from the body with a total body clearance of 10.3 ml/min.kg. If dosed as continuous drinking water medication (multiple dosing) steady-state concentrations of 0.5 mg (turkeys) to 0.8 mg (chicken) enrofloxacin per litre are achieved. A high mean volume of distribution (5 L/kg) indicated good tissue penetration of enrofloxacin. Concentrations in target tissues like lungs, liver, kidney, intestine and muscle tissue, exceed plasma concentrations by far. In poultry enrofloxacin is poorly metabolised to its active metabolite ciprofloxacin (approximately 5%). Enrofloxacin is eliminated from the body at a half-life of 6 hours. Protein binding in poultry is approximately 25%.

4. Indication

Treatment of infections caused by the following bacteria susceptible to enrofloxacin :

●Chickens

Mycoplasma gallisepticum, Mycoplasma synoviae, Avibacterium paragallinarum, Pasteurella multocida, Escherichia coli.

●Turkey

Mycoplasma gallisepticum, Mycoplasma synoviae, Pasteurella multocida, Escherichia coli.

Enrofloxacin should be used where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the active substance of choice.

5. Recommended Dosage / Mode of Administration

The product is administered via the drinking water.

Chickens and turkeys :

10 mg enrofloxacin/kg bodyweight per day for 3-5 consecutive days.

Treatment for 3-5 consecutive days; for 5 consecutive days in mixed infections and chronic progressive forms. If no clinical improvement is achieved within 2-3 days, alternative antimicrobial therapy should be considered based on susceptibility testing.

Via the drinking water. Always make sure that the entire dose offered has been consumed.

The medicated water should be made up fresh each day just before it is offered to the animals.

The drinking water must be medicated throughout the treatment period, and no other water source should be available.

Determine the bodyweight of the birds as accurately as possible in order to avoid underdosing.

Use only fresh pre-solutions, prepared every day before start of treatment. Pumping systems should be checked constantly to assure proper medication.

Empty the water system and fill it with medicated water before starting the treatment.

Calculate the daily quantity (ml) of Kenoxin 10% Oral Solution required for treatment period as follows :

Total number of birds x Average body weight in kg x 0.1= Total volume (ml) per day Kenoxin 10% Oral Solution may be put directly into the header tank or introduced via a water proportioner pump.

0000000
4603010
200401

6. Contraindications

Do not use for prophylaxis.

Do not use when resistance/ cross-resistance to (fluoro) quinolones is known to occur in the flock intended for treatment.

Do not use in the case of known hypersensitivity to the active substance, other (fluoro) quinolones or to any of the excipients.

7. Warning & Precautions

Special warnings for each target species

Treatment of Mycoplasma spp infections may not eradicate the organism.

Special precautions for use

i) Special precautions for use in animals

Official and local antimicrobial policies should be taken into account when the product is used.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

Since enrofloxacin was first authorised for use in poultry, there has been widespread reduction in susceptibility of E.coli to fluoroquinolones and emergence of resistant organisms.

Resistance has also been reported in Mycoplasma synovia in the EU.

Whenever possible, fluoroquinolones should only be used based on susceptibility testing.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance.

ii) Special precautions to be taken by the person administering the medicinal product to animals

Those with known hypersensitivity to (fluoro)quinolones should avoid contact with this product.

Avoid contact with skin and eyes.

Rinse any splashes from skin or eyes immediately with water.

Wash hands and exposed skin after use.

Do not eat, drink or smoke while using the product.

Keep out of children/ Jauhi ubat daripada kanak-kanak.

TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS ONLY

8. Interactions with Other Medicaments

In vitro, an antagonism was shown, when combining fluoroquinolones with bacteriostatic antimicrobial agents such as macrolides or tetracyclines and phenicols. The simultaneous application of substances containing aluminium or magnesium can impair the absorption of enrofloxacin.

9. Statement on usage during pregnancy and lactation

Do not use in laying hens producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

10. Adverse Effects/Undesirable Effects

None known

11. Overdose and Treatment

No adverse clinical symptoms were observed in chickens and turkeys treated respectively with doses up to 10 and 6 times higher than the therapy dose.

The use of fluoroquinolones during the growth phase combined with a marked and prolonged increase in the intake of drinking water, and hence active ingredient, possibly due to high temperatures, may potentially be associated with damage of the articular cartilage.

12. Withdrawal Period(s)

Chickens : Meat and offal : 7 days

Turkeys : Meat and offal : 13 days

Not authorised for use in birds producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

13. Storage Conditions

Store in a cool and dark place with dry condition.

Do not store above 30 °C .

14. Dosage Forms and packaging available

100 ml, 500ml and 1L

15. Shelf life (after first opening of container)

24 hours

16. Shelf life (after reconstitution or dilution)

24 hours

17. Name and Address of manufacturer/marketing authorization holder

Nutri Pharmax Sdn Bhd

6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.

18. Date of Revision of Package Insert : 01 April 2020



Manufactured by

KBNP, INC.

235-9 Chusaro, Sinam, Yesan, Chungnam, Korea



Distributed by

Nutri Pharmax Sdn.Bhd.

6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel : 04-5829377

1.사 이 즈	설명서 (100*162)
2.재 질	모조지
3.코 틱	코팅 없음
4.수 량	10,000매
5. Roll 폴림방향	
6.제작사	우 양
7.가 공	
※기존인쇄사양과 다르면 연락 주세요	