

OMS TIAMULIN 10% PREMIX POWDER

Composition

Each g contains:

Tiamulin hydrogen fumarate..... 100 mg

Product Description

OMS Tiamulin 10% Premix Powder is white or slightly yellowish powder.

Pharmacodynamics

Tiamulin is a bacteriostatic semi-synthetic antibiotic belonging to the pleuromutilin group of antibiotics and acts at the ribosomal level to inhibit bacterial protein synthesis.

Tiamulin has shown a in-vitro activity against a wide range of bacteria such as *Brachyspira hyodysenteriae*, *Brachyspira pilosicoli*, *Lawsonia intracellularis* and *Mycoplasma spp*.

Tiamulin is bacteriostatic at therapeutic concentrations and has been shown to act at the 70S ribosome level and the primary binding site is on the 50S subunit and possibly a secondary site where the 50S and 30S subunits join. It appears to inhibit microbial protein production by producing biochemical inactive initiation complexes, which prevent elongation of the polypeptide chain.

Mechanisms responsible for resistance development in *Brachyspira spp* to the pleuromutilin class of antibiotics are considered to be based on mutations at the ribosomal target site. Clinically relevant resistance to tiamulin requires combinations of mutations around the tiamulin binding site. Resistance to tiamulin may be associated with decreased susceptibility to other pleuromutilins.

Pharmacokinetics

Pig

Tiamulin is well absorbed in the pig following oral administration and widely distributed through the body.

Tiamulin has been shown to concentrate in the lung, a target tissue, and also in liver, where it is metabolised and excreted in the bile, the remainder is excreted via the kidney. Tiamulin which has not been absorbed or metabolised, passes down the intestines to the colon and concentrates there.

Chicken

Tiamulin is well absorbed in chickens after oral administration.

Tiamulin distributes widely through the body and to concentrate in the liver and kidney (sites of excretion) and in the lung. Excretion is mainly via the bile and kidney.

Indication

Pig

For the treatment of swine dysentery caused by *Brachyspira hyodysenteriae*.

For the treatment of colitis caused by *Brachyspira pilosicoli*.

For the treatment of ileitis caused by *Lawsonia intracellularis*.

For the treatment of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

Chicken

For the treatment of chronic respiratory disease (CRD) and air sacculitis caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*.

Mode of Administration

Orally. Mixed into feed.

Recommended Dosage

Swine:

Treatment of Swine Dysentery caused by *B. hyodysenteriae*, treatment of Porcine Colonic Spirochaetosis (colitis) caused by *B. pilosicoli*

Dosage: 5 – 10 mg tiamulin hydrogen fumarate/kg body weight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 – 200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Treatment of Porcine Proliferative Enteropathy (ileitis) caused by *L. intracellularis*

Dosage: 7.5 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 10 to 14 consecutive days. The dosage will normally be achieved by an inclusion level of 150 ppm tiamulin hydrogen fumarate in the finished feed providing that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.5 kg

Treatment of Enzootic Pneumonia caused by *M. hyopneumoniae*

Dosage: 5.0 – 10.0 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 – 200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Secondary infection by organisms such as *Pasteurella multocida* and *Actinobacillus pleuropneumoniae* may complicate enzootic pneumonia and require specific medication.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Chicken (broiler, laying and breeding hens):

Treatment of Chronic Respiratory Disease (CRD) caused by *M. gallisepticum* and air sacculitis and infectious synovitis caused by and *M. synoviae*.

Dosage – Treatment: 25 mg tiamulin hydrogen fumarate / kg bodyweight daily administered for the period of 3 to 5 consecutive days. This is normally achieved by an inclusion level of 250 – 500 ppm tiamulin hydrogen fumarate in finished feed provided that feed intake is unaffected. Inclusion levels in the higher range will in most cases be needed to avoid underdosing. In fast growing birds, e.g. broiler chickens during the first 2 – 4 weeks of life, inclusion levels in the lower range may be sufficient.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	2.5 – 5.0 kg

Contraindications

Animals should not receive products containing ionophores (monensin, narasin or salinomycin) during or for at least seven days before or after treatment with tiamulin. Severe growth depression or death may result.

Warnings and Precautions

Special warnings for each target species

In case of reduced feed intake, the inclusion levels in feed may need to be increased to achieve target dosage. Acute cases and severely diseased animals with reduced feed intake should be treated with a product of suitable formulation such as an injectable or water solution.

Special precautions for use in animals

It is sound clinical practice to base treatment on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of target bacteria.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

When mixing the veterinary medicinal product and handling the medicated feed, direct contact with eyes, skin and mucous membranes should be avoided. Personal protective equipment should be worn when mixing the veterinary medicinal product or handling the medicated feed: overalls, impervious gloves and either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator conforming to European Standard EN 140, with a filter to European Standard EN 143. Wash contaminated skin.

In case of accidental ingestion, seek medical advice immediately and show the package insert or label to the physician.

People with known hypersensitivity to tiamulin should administer the product with caution.

Interactions with Other Medicaments

Tiamulin has been shown to interact with ionophores such as monensin, salinomycin and narasin and may result in signs indistinguishable from an ionophore toxicosis. Animals should not receive products containing monensin, salinomycin or narasin during or at least 7 days before or after treatment with tiamulin. Severe growth depression, ataxia, paralysis or death may result.

If signs of an interaction do occur, administration of contaminated feed should be stopped immediately. The feed should be removed and replaced with fresh feed not containing the anticoccidials monensin, salinomycin or narasin.

Pregnancy and Lactation

Can be used in pigs during pregnancy and lactation.

Can be used in laying and breeding chickens.

Adverse Effects/Undesirable Effects

On rare occasions erythema or mild oedema of the skin may occur in pigs following the use of tiamulin.

Overdose and Treatment

Pigs: Single oral doses of 100 mg/kg bodyweight in pigs caused hyperpnoea and abdominal discomfort. At 150 mg/kg no CNS effects were noted except for sedation. At 55 mg/kg given for 14 days a transient salivation and slight gastric irritation occurred. A minimum lethal dose has not been established in the pig.

Poultry: The LD5 for chickens is 1290 mg/kg and turkeys 840 mg/kg bodyweight. The clinical signs of acute toxicity in chickens are – vocalization, clonic cramps and lateral recumbency. In turkeys signs of acute toxicity include clonic cramps, lateral or dorsal recumbency, salivation and ptosis. If signs of intoxication do occur promptly remove the medicated feed, replace with fresh unmedicated feed and apply supportive, symptomatic therapy.

Withdrawal Period(s)

Pigs

Treatment (at 5-10 mg/kg bwt): 6 days

Chickens

Meat and offal: 1 day

Eggs: 0 days

Storage condition

Store in a cool dry place, protected from light. Do not store above 30°C.
Keep out of reach of children/Jauhkan dari kanak-kanak.

Pack size(s)

Paper bag of 20kg

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)
Lot 16014(P.T. 24341), Jalan Nilam 3,
Bandar Nilai Utama,
71800 Nilai, Negeri Sembilan, Malaysia.
Tel: +606-799 9090 Fax:+606-799 9155

Date of revision of Package Insert

20/05/2024