



MYCOGAL 1.05 MIU/ml Solution for injection

COMPOSITION:

Name and strength of active substance:

Spiramycin adipate: 1.05 MIU/ml

Name and content of preservatives:

Benzyl alcohol: 0.1 ml/ml

PRODUCT DESCRIPTION:

It is a clear orange-coloured solution free from visible impurities.

PHARMACODYNAMICS:

It is a macrolide antibiotic with bacteriostatic action at usual doses and with bactericidal action at high doses that inhibits the biosynthesis of the bacterial proteins. Its spectrum of activity includes: Gram-positive bacteria (*Arcanobacterium pyogenes*, *Streptococcus dysgalactiae*, *Streptococcus agalactiae*, *Streptococcus uberis* and *Staphylococcus aureus*) and *Mycoplasma* spp.

PHARMACOKINETICS:

Peak plasma levels occur within 1-2 hours in most cases. It becomes widely distributed in tissues, and concentrations are higher than in plasma. Spiramycin tends to be concentrated in the spleen, liver, kidneys and particularly in the lungs. It is fixed to the plasmatic proteins in a proportion of 18-30%.

Spiramycin and its metabolites are excreted mainly in bile (>60%) and often undergo enterohepatic cycling. Urinary clearance is 4-20%.

In swine, spiramycin has a different behaviour in three phases: a first (absorption) in which the concentrations are increasing, a second one in which the concentrations decrease rapidly (after 4 hours) and a third one (excretion) from 6-8 hours in with the concentrations decrease slowly.

In bovine, spiramycin levels in plasma after intramuscular administration at doses of 25 mg/kg bw were high during the first 6 hours being observed a second peak between 1.5-6 hours after administration. Then, levels decrease steadily without disappearing at 120 hours with a half-life of elimination of 22-38 hours.

TARGET SPECIES: Cattle and swine.

INDICATION FOR USE:

Bovine: mastitis caused by *Arcanobacterium pyogenes*, *Streptococcus dysgalactiae*, *Streptococcus agalactiae*, *Streptococcus uberis* and *Staphylococcus aureus* spiramycin-sensitive; and respiratory infections caused by *Mycoplasma* spp. and *Arcanobacterium pyogenes* spiramycin-sensitive.

Swine: treatment of respiratory infections caused by *Mycoplasma* spp spiramycin-sensitive.

CONTRAINDICATIONS:

Do not use in case of hypersensitivity to the active substance, other macrolide antibiotics or to any of the excipients.

ROUTE, DOSAGE AND METHOD OF ADMINISTRATION:

Administration route: intramuscular.

25 mg of activity/kg bw (equivalent to 0.1 ml of MYCOGAL 1.05 MIU/mL/kg bw or equivalent to 1 ml of MYCOGAL 1.05 MIU/ML/10 kg bw) in single administration.

ADVICE ON CORRECT ADMINISTRATION:

The weight of animals must be determined as precisely as possible to avoid under-dosing.

Do not administer more than 15 ml in the same administration point.

SPECIAL WARNINGS:

Use in animals: Official and local antimicrobial policies should be taken into account when the product is used. Whenever possible, Spiramycin should be only used on susceptibility testing.

Use by the person during administration: People who are hypersensitive to macrolide antibiotic should avoid contact with the product. It should be handled with care to avoid self-injection or contact with the skin or the eyes. In case of contact, wash immediately with plenty of water. In the event of accidental self-injection consult your doctor and show the leaflet.

Use during pregnancy, lactation or lay: Studies in laboratory animals have not shown teratogenic effects, toxicity to foetus or mother. It has not been demonstrated the safety of the veterinary product during the gestation and/or lactation in target species. Use according to the evaluation benefit/risk assessment by the responsible veterinarian.

INTERACTIONS WITH OTHER MEDICAMENTS: None.

ADVERSE REACTIONS:

Allergic reactions in animals with hypersensitivity to macrolide antibiotics. Oedema and haemorrhage can occur in tissues near the injection site.

OVERDOSE AND TREATMENT: None known.

INCOMPATIBILITIES: None known.

WITHDRAWAL PERIOD:

Swine (meat): 22 days.

Bovine (meat): 36 days.

Milk: 11 days.

SPECIAL STORAGE PRECAUTIONS:

Store below 25°C. Store in a dry place and protect from light.

Shelf-life after first opening the immediate packing: 28 days.

Shelf-life before first opening the immediate packing: 4 years.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

PACKAGING MATERIAL AVAILABLE:

100 ml and 250 ml.

NAME AND ADDRESS OF MANUFACTURER:

Laboratorios Ovejero S.A. Ctra. León-Vilecha,30, 24192 León (SPAIN).

NAME AND ADDRESS OF PRODUCT REGISTRATION HOLDER:

Y.S.P. INDUSTRIES (M) SDN. BHD.

Lot 3, 5 & 7, Jalan P/7, Section 13,

Kawasan Perindustrian Bandar Baru Bangi,

43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679

To update the name and
address of PRH

DATE OF REVISION OF PACKAGE INSERT:

28 Nov 2018

To update the date
of revision

MAL 20162656HA

To update the MAL
number

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN

JAUHI DARIPADA KANAK-KANAK

FOR ANIMAL TREATMENT ONLY

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION