

T-LOX 66.8% W/W WATER SOLUBLE POWDER

T-Lox 66.8% w/w Water Soluble Powder appears to be a light yellow color powder.

Composition

Each g contains
Tylosin Tartrate 668 mg

Pharmacodynamics

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes Gram-positive bacteria including *Clostridium perfringens* and some Gram-negative strains such as *Pasteurella* and *Mycoplasma* spp. at concentrations of 16µg/ml or less.

Pharmacokinetics

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord.

Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

Indication

-For the control of *Mycoplasma synoviae* airsacculitis in chickens and *Mycoplasma gallisepticum* S6 in chickens and turkeys. In the field tylosin has also proved useful in reducing the level of infection following stress associated with live vaccination.

-As an aid in the control of outbreaks of necrotic enteritis in chickens caused by *Clostridium perfringens*.

-For the prevention and control of enzootic pneumonia, swine dysentery and other scour caused by organisms (e.g. *Lawsonia intracellularis*) sensitive to tylosin, in pigs.

-For the control of pneumonia in cattle associated with mycoplasmas and *Pasteurella multocida* sensitive to tylosin.

Recommended Dosage

Chickens and turkeys:

T-Lox 66.8% is administered in the drinking water at a concentration of 0.82 g per litre.

As an aid in the control of outbreaks of necrotic enteritis caused by *Clostridium perfringens* in chickens use T-Lox 66.8% in the drinking water for 5 days at a concentration of 0.25 g per litre water, to provide 20-50 mg/kg bw, depending on the age and the water consumption of the birds.

Pigs:

T-Lox 66.8% is administered in the drinking water to provide 25 mg tylosin/kg bodyweight. This may normally be achieved by adding 0.41 g per litre.

Calves:

1.65 g of T-Lox 66.8% per calf administered orally twice daily for seven to fourteen days. The tylosin tartrate may be incorporated into the milk or reconstituted milk replacer at the time of feeding.

Mode of Administration

Oral Administration.

Contraindication

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or to wildlife.

Do not use in known cases of hypersensitivity to tylosin or other macrolides.

Warning and Precautions

Special warnings for each target species

None.

Special precautions for use

Special precautions for use in animals

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator or a non-disposable respirator. Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interaction with Other Medicinal Product

None known.

Pregnancy and Lactation

No adverse effects to tylosin have been seen in fertility, multi-generation or teratology studies.

Adverse Effects / Undesirable Effects

None known.

Overdose and Treatment:

There is no evidence of tylosin toxicity in animals, at dose rates of up to 1000 mg/kg.

Withdrawal Period (s)

Chickens: For the control of chronic respiratory disease: Meat: 1 day, Egg: 0 day

As an aid in the control of outbreaks of necrotic enteritis caused by *Clostridium perfringens*: Meat: 0 day, Egg: 0 day

Turkeys: For the control of chronic respiratory disease: Meat: 0 day

Pigs: For the prevention and control of enzootic pneumonia, swine dysentery and other scour caused by organisms (e.g. *Lawsonia intracellularis*) sensitive to tylosin: Meat: 0 day

Calves: For the control of pneumonia associated with mycoplasmas and *Pasteurella multocida* sensitive to tylosin: Meat: 14 days

Storage Conditions

Store below 30°C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children / Jauhkan ubat dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf-life after opening: 7 days

Shelf-life after reconstitution: 24 hours

Packaging Available

200g, 500 g & 1 kg.

Product Holder/ Manufacturer

NAM PHARMA SDN BHD

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Date of revision

09-09-2024