

CHEMIX[®]

ORAL SOLUTION 200MG/ML

Ingredient(s):

Each ml contains:

Flumequine	200mg
Benzyl Alcohol (as solvent)	0.02ml

Pharmacodynamics:

Flumequine is an antibacterial agent belonging to the group fluoroquinolone bactericidal.

It acts on bacterial DNA, blocking its replication by inhibiting the enzyme subunit of DNA gyrase, the enzyme is involved in the formation of the DNA helix. It is effective on gram-negative bacteria. Flumequine resistance occurs by the alteration of DNA gyrase (topoisomerase II) by mutating the Gyr-A subunit and less frequently by mutating the ParC (Topoisomerase IV) subunit.

Other resistance mechanisms appear when the bacteria decrease the permeability of the membrane, thus preventing the passage of antimicrobial or increasing the active compound transport out of the cell.

Pharmacokinetics:

After oral administration, 10% of the dose is absorbed, reaching peak plasma concentration at 2 hours.

It is widely distributed in tissues. Its plasma protein binding is 74.5% in blood, the unbound flumequine presents in the form of hydroxylated metabolites with low activity. Approximately 40-60% of the absorbed dose is excreted in the urine, and the remainder is excreted as inactive glucuronide metabolite. A small portion is excreted in faeces.

Indication:

The product is indicated for the treatment of infections caused by bacteria sensitive to flumequine:

Calves: bronchopneumonia, enteritis colibacillosis, neonatal diseases, salmonellosis

Swine: colibacillosis, enteritis, gastroenteritis, neonatal diseases

Sheep: colibacillosis, septicemia, pasteurellosis

Poultry (broilers and turkeys): colibacillosis, salmonellosis, pasteurellosis

Target Species:

Calves, swine, sheep and poultry (broilers and turkeys).

Recommended Dose:

Calves: 5-10 mg/ kg body weight, equivalent to 1.25-2.5 ml product / 50 kg body weight. Dose is repeated every 12 hours for 5 consecutive days.

Adult swine: 6 mg/ kg body weight, equivalent to 3 ml of product / 100 kg body weight, twice a day at an interval of 12 hours for 3-5 consecutive days.

Piglet: 12 mg/ kg body weight, equivalent to 3 ml of product / 50 kg body weight, the dosage is repeated every 12 hours for 3-5 consecutive days.

Sheep: 6 mg / kg bodyweight every 12 hours (equivalent to 1.5 ml / 50 kg body weight), for 4 – 6 consecutive days.

Poultry (chicken and turkeys): 12-24 mg/ kg body weight, equivalent to 1 ml product / 8-16 kg body weight, for 3-5 consecutive days.

Mode of Administration

Oral administration via drinking water.

Contraindication

Never use in animals suffering from hepatic or renal impairment. Do not use in case of hypersensitivity to the active substance or excipients.

Do not use in animals producing milk for human consumption.

Do not use in birds producing eggs for human consumption.

Warning and Precautions

Special precautions for use in animals

1. Do not expose treated animals to sunlight because of the risk of photosensitivity.
2. When using the product should take official and local policies on antimicrobial products into account.
3. Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly or are expected to respond poorly to other classes of antimicrobials. Fluoroquinolones should be used only after performing susceptibility testing. Use of the product deviating from the instructions given in the insert may increase the prevalence of bacteria resistant to quinolones and may decrease the effectiveness of treatment with other fluoroquinolones due to the potential for cross-resistance.

Special precautions to be taken by the person administering the product to animals

1. Flumequine may cause hypersensitivity (allergy) following inhalation, ingestion or skin contact. There may be cross-hypersensitivity with other quinolones.
2. People with known hypersensitivity to quinolone should avoid contact with the drug.
3. Manipulate the product with caution. During handling, wear personal protective equipment consisting of gloves, overalls, safety glasses.
4. Avoid direct contact with skin and mucous membranes. In case of accidental contact, wash immediately with plenty of water. If symptoms appear after exposure, such as a rash, consult a doctor.
5. In case of accidental ingestion, seek medical advice immediately and show the package insert or label. Swelling of the face, lips and eyes are more severe signs that require urgent medical attention.
6. No smoking, eating or drinking while medicine is handled.

Interaction with Other Medicaments

Do not administer concurrently with trimethoprim.

Pregnancy and Lactation

It has not been demonstrated the safety of the product during pregnancy and lactation in sows. Use only according to the benefit / risk assessment by the responsible veterinarian.

Do not use in laying birds or within 4 weeks before the start of the laying period.

Side Effects

There may be occasional digestive disorders (vomiting or diarrhea).

May occur nervous hyperexcitability (rarely) or photosensitivity (skin allergies).

Environmental Property

Non-hazardous.

Symptoms and Treatment of Overdose

May experience digestive or nervous disorders, but they decrease in intensity when treatment is stopped.

Storage Condition

Store in a cool place below temperature 30°C. Protect from direct sunlight.

Shelf Life

3 years.

Withdrawal Periods

Meat:

Calves and sheep: 10 days

Swine: 2 days

Poultry: (Broilers) 2 days; (Turkeys): 10 days

Maximum Residual Limit (MRL)

Bovine, Swine, Sheep and Poultry:

Muscle: 50 µg/kg

Liver: 100 µg/kg

Kidney: 300 µg/kg

Product Description

A yellowish to brownish yellow clear solution.

Plastic Bottle 1 L & 5 L

To declare Y.S.P. INDUSTRIES VIETNAM CO., LTD.
as Manufacturer and Y.S.P. INDUSTRIES (M) SDN.
BHD. as Product Registration Holder

**Made by YSP**

Manufacturer:

Y.S.P. INDUSTRIES VIETNAM CO., LTD

Road No.6, Nhon Trach 3 Industrial zone, Nhon Trach,
Dong Nai province, Vietnam Tel: +8461 3569 993

Product Registration Holder :

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679

To update the Date of
revision

Date of revision: 07 Sep 17