

PACKAGE INSERT

Brand or product name	Tylolide Granules for Oral Solution 100 % w/w
Name and Strength of Active Substance/ Preservative(s)	Composition per tub: Tylosin (as tartrate).....100,000,000 IU (Equivalent to 100 g of tylosin per tub or equivalent to 110 g of tylosin tartrate per tub)
Product Description	Almost white to light yellow granules
Pharmacodynamics	Tylosin is a macrolide antibiotic produced by a strain of <i>Streptomyces fradiae</i> . It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms. The tylosin spectrum of activity includes amongst others Gram-positive bacteria, some Gram – negative strains such as <i>Pasteurella</i> , and <i>Mycoplasma spp.</i>
Pharmacokinetics	In most species peak plasma concentrations have been attained 1 to 2 hours after administration of tylosin. Compared to plasma levels clearly higher tissue concentrations have been observed. Tylosin was extensively metabolised.
Environmental Properties	Most of the residues are excreted in faeces predominantly consisting of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.
Indication	<u>Calves</u> : Treatment and prevention of pneumonia caused by <i>Mycoplasma spp</i> when the disease has been established in the herd. <u>Pigs</u> : Treatment and prevention of enzootic pneumonia caused by <i>Mycoplasma hyopneumoniae</i> and <i>Mycoplasma hyorhinis</i> when the disease has been established in the herd. Treatment and prevention of Porcine Intestinal Adenomatosis (Ileitis) associated with <i>Lawsonia intracellularis</i> when the disease has been established in the herd. <u>Chickens</u> : Treatment and prevention of chronic respiratory diseases (CRD) caused by <i>Mycoplasma gallisepticum</i> and <i>Mycoplasma synoviae</i> when the disease has been established in the flock. Treatment and prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> when the disease has been established in the flock. <u>Turkeys</u> : Treatment and prevention of infectious sinusitis caused by <i>Mycoplasma gallisepticum</i> when the disease has been established in the flock.
Recommended Dosage	1.1 gram of the veterinary medicinal product corresponds to 1 gram of tylosin. The dosages are as follows:

Calves:

10 – 20 mg tylosin per kg BW (corresponding to 11 – 22 mg of the veterinary medicinal product per kg BW), twice daily (corresponding to a daily dose of 20 – 40 mg tylosin per kg BW), for 7 - 14 days.

Turkeys:

75 – 100 mg tylosin per kg BW per day (corresponding to 82.5 – 110 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

Chickens:

For the treatment of chronic respiratory disease:

75 – 100 mg tylosin per kg BW per day (corresponding to 82.5 – 110 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

For the treatment of necrotic enteritis:

20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product) for 3 days.

Pigs:

For the treatment of enzootic pneumonia:

20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product per kg BW) for 10 days.

For the treatment of ileitis or PIA:

5 – 10 mg tylosin per kg BW per day (corresponding to 5.5 - 11 mg of the veterinary medicinal product per kg BW) for 7 days.

For the preparation of the medicated water/milk/milk-replacer the body weight of the animals to be treated and their actual daily water/milk/milk-replacer consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system. To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{..... mg tylosin per kg bodyweight per day} \times \text{Average bodyweight (kg) of the animals to be treated}}{\text{Average amount of drinking water or milk / animal (l)}} = \text{.....mg tylosin / l of drinking water}$$

Sufficient access to the system of water supply should be available for the animals to be treated to ensure adequate water consumption. No other source of drinking water should be available during the medication period.

Should there be no clear response to treatment within 3 days the treatment approach should be reconsidered. After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance which might support development of resistance.

	Medicated water, milk or milk replacer should be replaced every 24 hours. If individual animals show signs of a serious infection such as a reduced water or feed intake, then they should be treated individually, such as by injection.
Mode of Administration	Orally administration dissolved in drinking water or milk.
Contra-indications	Do not use in known cases of hypersensitivity to tylosin or other macrolides.
Warnings and Precautions	a) Special warnings for each target species Due to likely variability (time, geographical) in susceptibility of bacteria to tylosin, bacteriological sampling and susceptibility testing are recommended. Under-dosing and treating for an insufficient length of time are considered to promote the development of resistance in bacteria and should be avoided. b) Special precautions for use i) Special precaution for use in animals: Animals with acute infections may have a reduced water and feed consumption and should be treated with a suitable injectable veterinary medicinal product first. Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or wildlife. ii) Special precautions for the person administering the veterinary medicinal product to animals Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Wash hands after use. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Do not handle the product if you are allergic to ingredients in the product.

	If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.
Interactions with Other Medicaments	Lincosamides and aminoglycoside antibiotics antagonise the activity of tylosin.
Statement on usage during pregnancy and lactation	Laboratory studies in mice and rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.
Adverse Effects/Undesirable Effects	In pigs, adverse reactions have been observed, including diarrhoea, pruritus, erythema of the skin, swelling of the vulva, rectal edema and prolapse. These reversible signs appeared 48-72 hours after start of treatment.
Overdose and Treatment	There is no evidence of tylosin toxicity in rats, at dose rates of up to 1000 mg/kg by the oral route. There is no evidence of tylosin toxicity in chickens, turkeys, pigs or calves when administered orally at up to three times the recommended dose.
MRL	<u>Cattle:-</u> Muscle, liver, kidney, fat – 0.2 ppm Milk – 100 µg/kg (0.1 ppm) <u>Pig:-</u> Muscle, liver, kidney, Skin and fat – 0.2 ppm <u>Chickens:-</u> Muscle, liver, kidney, skin and fat – 0.2 ppm <u>Turkeys:-</u> Muscle, liver, kidney, skin and fat – 0.2 ppm Eggs - 300 µg/kg (0.3 ppm)
Incompatibilities(for injections only)	None known.
Withdrawal period(s)	Meat and offal: - Poultry: 1 day. - Turkeys: 2 days. - Pigs: 1 day. - Calves: 12 days. Eggs: zero day.
Storage conditions (may be omitted if the information is stated on	Storage at 25 °C/60 % R.H.

the label or outer carton labels)/(should be supported by stability data)	
Dosage forms and packaging available	Dosage form: Granules for Oral Solution Packaging available: Tub of 100
Name and address of manufacturer	Laboratoires biové 3, Rue De Lorraine 62510 ARQUES, France
Name and Address registration holder	Weissen Company (M) Sdn Bhd Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia
Date of revision of Package insert	01.07.2024
Shelf-life of the veterinary product as packaged for sale	36 months
Shelf-life after first dilution or reconstitution according the directions (where relevant)	24 hours after dissolution in water at 20 g/L 24 hours after dissolution in water at 0.1 g/L 2 hours after dissolution in reconstituted milk at 0.5 g/L