

NEOCIN 49% POWDER

Description and Composition

NEOCIN 49% POWDER is a white to yellow white powder. Each kg contains 490g Neomycin (as sulphate).

Pharmacodynamics

Neomycin is an aminoglycoside antibiotic. Its spectrum of action includes many gram-negative organisms including *Escherichia coli* and enterobacteria.

The aminoglycosides bind to the 30S ribosomal subunit of the bacterium. By binding to the ribosome, aminoglycosides inhibit the translocation of the tRNA during translation and therefore inhibit the synthesis of proteins required for the growth of the bacterium.

Pharmacokinetics

Neomycin is poorly absorbed from the gastrointestinal tract.

Neomycin is almost exclusively eliminated in the faeces after oral administration.

Indication

For the treatment of gastrointestinal infections caused by bacteria sensitive to neomycin in piglets and poultry.

Recommended Dose

Piglets: 2.04g to 3.06g of Neocin 49% Powder per 100kg body weight (equivalent to 10mg to 15mg of Neomycin per kg body weight) every 12 hours for 3 to 5 consecutive days.

Poultry: 4.08g to 6.12g of Neocin 49% Powder per 10 litres of drinking water (equivalent to 20mg to 30mg of Neomycin per kg body weight) for 3 to 5 consecutive days, to be adjusted according to the actual consumption of the animals based on the dosage weight (mg/kg).

Route of Administration

For oral administration via drinking water.

Contraindication

Do not use in animals known to be allergic to aminoglycosides.

Warning and Precautions

Special warnings for each target species

None.

Special precautions

i) Special precautions for use in animals

This soluble oral powder is intended to target species indicated in the "Indications for use".

This powder for oral solution is intended to be dispersed in water.

Avoid administration in case of severe renal impairment.

At low concentrations, all aminoglycosides can only be bacteriostatic.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

People with known hypersensitivity to aminoglycosides should avoid contact with the veterinary medicinal product.

In case of appearance after exposure, symptoms such as rash, consult a doctor and show this cautionary statement.

Swelling of the face, lips or eyelids or difficulty breathing are more serious symptoms and require care.

Other precautions

None.

Interaction with Other Medicaments

Some aminoglycosides are more efficiently transported at the cell membrane than others, and thus have a higher antibacterial activity. Synergy is common when aminoglycosides and β -lactam antibiotics (penicillins and cephalosporins)

are used in combination. The lesion of the cell wall induced β -lactam compounds allows high incorporation of aminoglycosides by the bacterial cell due to greater accessibility to the cell membrane. Renal toxicity may become evident in the coadministration of aminoglycoside and other potentially toxic agents (methoxyflurane, amphotericin B, cisplatin and some cephalosporins).

Pregnancy and Lactation

Studies on laboratory animals have not demonstrated any evidence of teratogenic effects of neomycin. The safety of the product has not been evaluated in the target species. Use of the drug will be based on the evaluation of the benefit / risk carried out by the veterinarian.

Side Effects

Classically aminoglycosides are poorly absorbed (less than 10%) in the gastrointestinal tract. However enteritis and other pathological changes can enable higher absorption resulting in impaired renal function, increasing toxicity levels causing acute tubular necrosis with secondary interstitial injury.

Symptoms and Treatment of Overdose

Non-oliguric renal failure is the usual clinical situation during aminoglycosides intoxication; it is generally reversible although healing may be delayed. Monitor renal function during treatment; polyuria, decreased urinary osmolality, enzymuria, proteinuria and cylindruria increase in sodium excretion fraction are signs of nephrotoxicity of aminoglycosides.

Storage Condition

Store below 30°C.

Shelf life

3 years. To be used within 24 hours after dilution. After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Period

Piglets: Meat and offal: 14 days

Poultry:

Meat and offal: 14 days

Eggs: Zero days

Packing

100g, 500g and 25kg

Revised date: 17.11.2016

	Manufactured by:
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 info@rangepharma.com.my