

FOSVET-T Water Soluble Powder

FOR ANIMAL USE ONLY

Brand or Product Name :

FOSVET-T Water Soluble Powder

Name and Strength of Active Substance(s)

Each gram of FOSVET-T Water Soluble Powder contains Fosfomycin Calcium 200mg and Tylosin Tartrate 50mg.

Product Description

FOSVET-T Water Soluble Powder is a white or light milky white powder.

Pharmacodynamics and Pharmacokinetics Properties

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms. The tylosin spectrum of activity includes Gram-positive bacteria including *Clostridium perfringens* and some Gram-negative strains such as *Pasteurella* and *Mycoplasma* spp. at concentrations of 16µg/ml or less. Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose. Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord. Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

Fosfomycin is a phosphoenol pyruvate analog that irreversibly inhibits pyruvyl transferase, the enzyme that catalyzing the first step of peptidoglycan biosynthesis. Fosfomycin is a broad-spectrum antibiotic produced by various *Streptomyces* spp. It is effective against many Gram-positive and Gram-negative. It is particularly active against many Enterobacteriaceae, including *E.coli* and also is active against *Pasteurella*, *Salmonella*, and *Staphylococcus*. After PO administration, absorption of fosfomycin occurs throughout the digestive tract. Fosfomycin has no metabolic transformation and is excreted in active form mostly through renal elimination.

Indication

For use in poultry (broilers; light and heavy breeders) in the treatment of bacterial infections caused by Gram-positive and Gram-negative bacteria including *E.coli*, *Salmonella*, *Pasteurella*, *Staphylococcus* and *Streptococcus*, and *Mycoplasma* spp. such as *Mycoplasma gallisepticum* and *Mycoplasma paragallinarum* that are sensitive to Fosfomycin and Tylosin.

Recommended Dosage

Administer orally by mixing in drinking water.

Poultry (*Broilers; light and heavy breeders*): Use 160mg of FOSVET-T per kilogram of body weight for 3 to 7 days, as instructed by veterinarians.

Mode of Administration

Administer orally by mixing in drinking water. Medicated water should be made up immediately prior to provision. Medication of the water supply should be continuous during the treatment period and no other source of water should be available. The product should be given over a minimum period of 6 hours per day to allow all poultry to drink.

Contraindications

None.

Warnings and Precautions

Discontinue treatment 7 days before slaughter.

Keep out of reach of children.

Do not administer to fowls producing eggs for human consumption.

Interactions with Other Medicaments

No significant drug interaction is reported.

Statement on usage during pregnancy and lactation

Safety data not available. Use with caution.

Adverse Effects / Undesirable Effects

None.

Overdose and Treatment

None.

Withdrawal Period(s)

7 days.

Shelf-life

Shelf life as packaged in original container: 3 years from the date of manufactured.

Shelf life after immediate opening of the container: 24 hours

Shelf life after reconstitution in drinking water: 24 hours

Allowable Maximum Residual Limit (MRL)

Tylosin : Muscle, liver, kidney, fat, egg = 200 µg/kg

Fosfomycin : Muscle, fat, liver, kidney and edible offal = 500µg/kg.

Storage Conditions

Store in a dry place below 25°C. Keep out of reach of children. Any opened sachet or unused medicated water to be discarded after 24 hours.

Dosage Forms and Packaging Available

Available in 100g, 200g, 500g and 1000g powder per sachet, pack in 10, 20, 50 and 100 sachets per box.

Disposal of Container

Traces of medicated powder left in an opened aluminum sachet can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned aluminum sachet can be safely disposed in trash according to local government regulations.

Name and Address of Manufacturer and Registration Holder

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2016-11-19