

OMS LINCOCEN 4.4% W/W PREMIX POWDER

Composition:

Each gm contains:

Lincomycin (as lincomycin hydrochloride).....44mg

Product Description:

OMS Lincocen 4.4%w/w Premix Powder is an off white powder.

Pharmacodynamics:

Lincomycin is a lincosamide antibiotic and is active against a wide range of Gram-positive aerobic and anaerobic bacteria, anaerobic Gram-negative bacteria and mycoplasma. Lincosamides are generally bacteriostatic. It is active against the primary causal agent of swine dysentery, *Serpulina hyodysenteriae* and the exacerbating anaerobes, *Bacteroides*, *Fusobacterium* and *Clostridium* spp, and also of mycoplasmal (enzootic) pneumonia in pigs, *Mycoplasma hyopneumoniae*.

Pharmacokinetics:

Lincomycin has approximately 50% systemic bioavailability from oral administration in pigs and is significantly re-circulated via the biliary route.

Indications:

Broilers

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

Swine

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*. For reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

Mode of Administration:

Orally. Mixed into feed.

Recommended Dosage:

To be administered orally via the feed.

Broilers:

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers

2 grams per ton of feed.

Swine:

For treatment of swine dysentery

100 grams per ton of feed, feed as sole ration for 3 weeks or until signs of disease disappear.

For control of swine dysentery

40 grams per ton of feed, feed as sole ration, for use in swine on premises with a history of swine dysentery but where symptoms have not yet occurred, or following use of lincomycin at 100 grams / ton for treatment of swine dysentery.

For control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*.
100 grams per ton of feed, feed as sole ration for 3 weeks or until signs of disease disappear.

For reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.
200 grams per ton of feed, feed as sole ration for 3 weeks.

Contraindications:

Do not administer to animals with known hypersensitivity to the active ingredient.
Not for use in animals other than the target species. Lincomycin may cause fatal enterocolitis in horses, rodents and rabbits.

Warnings and Precautions:

Special warnings for each target species
None.

Special precautions for use in animals

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Persons known to be hypersensitive (allergic) to lincomycin and/or soybean millfeed should not handle this product.

Care should be taken not to inhale any dust. The wearing of approved dust masks (either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143) and safety glasses is recommended during the handling and mixing of this product.

Skin contact should be avoided.

Wash hands and any exposed skin with soap and water immediately after use.

Interactions with Other Medicaments:

Clinical incompatibility may exist between lincomycin and erythromycin due to competitive binding at the ribosomal site of action.

Pregnancy and Lactation:

Not applicable.

Adverse Effects/Undesirable Effects:

May occasionally cause transient soft stools and/or mild swelling of the anus within the first two days of treatment. Very rarely some pigs may show reddening of skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing lincomycin treatment.

Overdose and Treatment:

Not applicable.

Withdrawal Period(s):

Swine:

No drug withdrawal period is required before slaughter of swine fed OMS LINCOCEN 4.4% W/W PREMIX POWDER.

Broilers:

No drug withdrawal period is required before slaughter of birds fed OMS LINCOCEN 4.4% W/W PREMIX POWDER.

Not for use in layers and breeders.

Storage conditions:

Store in a cool dry place, protected from light. Do not store above 30°C.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Pack size(s):

Paper bag of 10kg.

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)

Lot 16014(P.T. 24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

Tel: +606-799 9090 Fax:+606-799 9155

Date of revision of Package Insert:

~~11/01/2017~~ 12/07/2023