

CLEAROGOUT[®]

Alkalinising Effervescent Granules

COMPOSITION:

Each 4g sachet contains:

Sodium bicarbonate.....	1,760 mg
Sodium citrate anhydrous	630 mg
Citric acid anhydrous.....	720 mg
Tartaric acid.....	890 mg

Also contains lime flavour.

DESCRIPTION:

Appearance of powder: White to off-white granular powder with odour of lime flavour.

Appearance of solution: Clear to slight haze solution with odour of lime flavour.

PHARMACOLOGY:

Sodium bicarbonate acts as a systemic and urinary alkaliniser by increasing the excretion of free bicarbonate ions in the urine, thus effectively raising the urinary pH. By maintaining alkaline urine, the actual dissolution of uric acid stones may be accomplished. It is also a valuable adjuvant when administered with uricosuric agents in gout therapy, since urates tend to crystallise out of acidic urine.

Sodium Citrate and Citric Acid:

A rise in urinary pH increases the solubility of cysteine in the urine and the ionisation of uric acids to more soluble urate ion. By maintaining alkaline urine, the actual dissolution of uric acid stones may be accomplished.

Tartaric acid in combination with bicarbonates act as the acid component of effervescent granules.

PHARMACOKINETIC:

Sodium citrate is metabolised to bicarbonates, which increases urinary pH by increasing the excretion of free bicarbonate ions, without producing systemic alkalosis when administered in recommended doses. Sodium bicarbonate is excreted through renal and also via lung by forming CO₂. Sodium citrate, citric acid and absorbed tartaric acid are excreted through urine.

INDICATIONS:

Prevention of uric acid crystallisation in gout therapy & uric acid stones (kidney stones).

For relieving discomfort in mild urinary tract infections and symptomatic relief of painful dysuria. Enhance the action of certain antibiotics, especially sulphonamides.

DOSAGE AND ADMINISTRATION:


4g to 8g (1 to 2 sachets) dissolved in cold water four times daily or as recommended by a pharmacist or doctor.

ROUTE OF ADMINISTRATION:

Oral administration.

CONTRAINDICATIONS:

- Renal failure or hypernatremia.
- Concurrent hexamine mandelate or hexamine hippurate therapy.
- Caution is advised in overt and occult cardiac failure.
- Concomitant use of urinary alkalinisers and quinolone antibiotics should also be avoided because crystalluria may be more likely to occur in alkaline urine.



PRECAUTIONS/WARNINGS:

Those on a low sodium diet should take into account of the sodium content (650 mg per sachet) of this product.

DRUG INTERACTIONS:

Alkalinisation of the urine due to the use of CLEAROGOUT, theoretically, may result in a decreased therapeutic effect of the following medications: chlorpropamide, lithium, salicylates and tetracyclines. Alternatively, alkalinisation of the urine due to the use of CLEAROGOUT, theoretically, may result in an increased therapeutic effect of the following medications: amphetamines and ephedrine/pseudoephedrine.

Antacid: Concurrent use of antacids with sodium citrate and sodium bicarbonate may promote the development of calcium stones in patients with uric acid stones and may also cause hypernatremia. Concurrent use of aluminium-containing antacids with salts can increase aluminium absorption, possibly resulting in acute aluminium toxicity, especially in patients with renal insufficiency.

Quinolones: Citrates may reduce the solubility of ciprofloxacin, norfloxacin or ofloxacin in the urine. Patients should be observed for signs of crystalluria and nephrotoxicity.

Laxatives: Concurrent administration of citrates with laxatives may have an additive effect.

Hexamine hippurate/mandelate (inactive) is hydrolysed to formaldehyde (active) in acidic urine. Urinary alkalinisation prevents the formation of formaldehyde and thus reduces the antibacterial effect of hexamine.

PREGNANCY:

Effects of this product on pregnant women have not been carried out.

NURSING MOTHERS:

Caution should be exercised when administered to a nursing mother.

SIDE EFFECTS/ADVERSE REACTIONS:

The tartrate component of CLEAROGOUT may be incompletely absorbed. Because of this CLEAROGOUT may exert a mild laxative effect. Prolonged and excessive use may cause a systemic alkalosis and/or hypernatremia.

OVERDOSAGE:

Overdosage may result in metabolic alkalosis. CLEAROGOUT should be discontinued, appropriate therapy instituted and electrolyte and acid-base determinations should be carried out.

PACK SIZE:

Effervescent granules, 4g per sachet. 1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20, 24, 25, 28, 30, 40, 60, 80, 90, 100, 120, 240 sachets per pack.

PRODUCT REGISTRATION HOLDER:

Pahang Pharmacy Sdn. Bhd. Registration No. 197101000448 (10826-K)
Lot 5979, Jalan Teratai, 5½ Miles off Jalan Meru, 41050 Klang, Selangor.
INFOline: 603-6142 6570 (Mon-Fri; 9am-5pm)
Email: info@livewell2u.com

MANUFACTURER:

Noripharma Sdn. Bhd. Registration No. 200701034604 (792633-A)
Lot 5030, Jalan Teratai, 5 ½ Miles off Jalan Meru, 41050 Klang, Selangor.

DATE OF REVISION: 29 Jan 2024