

PVG Salinomycin 12% (w/w) Granule Premix

COMPOSITION

Salinomycin Sodium (120mg/g of product) equivalent to Salinomycin (116.6mg/g of product).

PRODUCT DESCRIPTION

Almost white to brown granules

PHARMACODYNAMICS

The coccidiocidal effect of salinomycin is directed mainly against free merozoites and against first and second generation schizont. Like all other polyene antibiotics, salinomycin acts as an ionophore and uses this property to facilitate the influx of cations into the cell. Salinomycin transports sodium and potassium ions into the cell in exchange for protons. This leads to a breakdown of the natural transmembrane ion gradient, a pH shift and the entry of water into the cell. The associated increase in pressure causes a destruction of intercellular structures and thus cell death. Salinomycin significantly reduces lesions caused by coccidia and improves body weight gain and feed tolerance compared to untreated animals.

PHARMACOKINETICS

Salinomycin is rapidly distributed and eliminated. The serum concentration kinetics correspond most closely to a two-compartment model with a distribution phase that ends after one hour. Since the drug is lipophilic and can easily diffuse through cell membranes, the distribution of salinomycin in tissue is greater than in serum. The highest concentrations are found in the liver, followed by kidneys, muscles, fat, heart and skin. The drug can no longer be detected in tissue after 48 hours, except in the liver, where traces can be measured up to 72 hours after administration.

INDICATIONS

For the prevention of *coccidiosis* caused by *Eimeria tenella*, *E.necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* in broiler and layer replacement chickens and pullets.

RECOMMENDED DOSAGE

Chickens: Thoroughly mix 500g of PVG SALINOMYCIN 12% (W/W) GRANULE PREMIX per one tonne of finished feed.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

Other anticoccidial drugs should not be added to feeds containing this product.

WARNING AND PRECAUTIONS

This product should not be used concurrently with Tiamulin/other antibiotics. The recommended inclusion dose rates should not be exceeded. This product must not be fed to horses or any other animal or bird not included on the label claim.

INTERACTIONS WITH OTHER MEDICAMENTS

Other anticoccidial drugs should not be added to feeds containing this product. This product should not be used concurrently with Tiamulin antibiotics.

PREGNANCY AND LACTATION

DO NOT USE in laying hens producing eggs for human consumption or within 7 days of laying where eggs or egg products are to be used for human consumption or processing. DO NOT USE in animals producing milk for human consumption.

SIDE EFFECTS

Not known.

OVERDOSE AND TREATMENT

The recommended inclusion dose rates should not be exceeded.

WITHDRAWAL PERIOD

MEAT: NIL

MILK: DO NOT USE in animals producing milk for human consumption.

EGGS: DO NOT USE in birds laying eggs for human consumption or within 7 days where eggs or egg products are to be used for human consumption or processing.

PRODUCT REGISTRATION NO.: MALXXXXX

IMPORTER & PRODUCT REGISTRATION HOLDER

QL PACIFIC VET GROUP SDN BHD

NO.886-C8, JALAN SUBANG 9, TAMAN PERINDUSTRIAN SUBANG, 47600 SUBANG JAYA, SELANGOR

MANUFACTURER

SHANDONG LUKANG SHELILE PHARMACEUTICAL CO., LTD.

50 HUALU ROAD, ZOUCHEG INDUSTRIAL PARK, ZOUCHEG, SHANDONG, P. R. CHINA

PACK SIZE

25KG

BATCH NO.: XXXXXX

MANUFACTURING DATE: XXXXXX

EXPIRY DATE: XXXXXX

DO NOT STORE ABOVE 30°C

JAUHI UBAT DARI KANAK-KANAK / KEEP OUT OF REACH OF CHILDREN

UBAT TERKAWAL / CONTROLLED MEDICINE

FOR VETERINARY USE ONLY

Date of Revision: 24.10.2025