

Veterinary Package Insert

TP-Apra 500 WSP Apramycin Sulphate 500 mg/g Water Soluble Powder

PRODUCT DESCRIPTION

A light yellow or light brown colour powder, which contains 500 mg/g of apramycin sulphate in 225g or 450g sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Apramycin is a broad-spectrum aminocyclitol antibiotic produced by a strain of *Streptomyces tenebrarius*. Apramycin is bactericidal at minimum inhibitory concentrations. Apramycin is effective against both Gram-negative and Gram-positive bacteria. It is effective against most field strains of *Escherichia coli* and salmonellae.

The oral administration of apramycin is intended for antimicrobial activity within the gut; apramycin is poorly absorbed, particularly in older animals:

INDICATION

Treatment of bacterial enteritis associated with organisms susceptible to apramycin in pigs, colibacillosis and salmonellosis in calves *Escherichia coli* septicaemia in young chickens.

RECOMMENDED DOSAGE AND ADMINISTRATION

The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder has to be adjusted accordingly.

Add the appropriate quantity of the product to 5 – 10 times its own volume of water. Stir well and allow the solution to stand for a few minutes and then stir again. When completely dissolved, add the full volume of drinking water and stir well.

For treatment in pigs, TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder should be administered via the drinking water. Treated pigs should consume sufficient medicated drinking water to obtain 7.5 – 12.5 mg apramycin activity per kg of bodyweight daily for seven days. This may be achieved by adding 2.4g of TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder per 20 litres of water.

As a guide, average daily water consumption for fattening pigs is shown in the following table:

Bodyweight (kg)	Average Consumption (litres)
5	1.10
10	1.75

For treatment in calves, TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder should be administered in the drinking water, milk or milk replacer to receive 20 – 40 mg apramycin activity per kg of bodyweight daily according to the severity of the disease. Continue treatment for five days. The full dose should be given once daily according to the following table:

Total Bodyweight to be treated	TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder	Apramycin (activity per day)
50 kg	2.4 – 4.7 g	1.0 to 2.0 g
250 kg	11.8 – 23.6 g	5.0 to 10.0 g
2500 kg	118.2 – 236.4 g	50.0 to 100.0 g

Solutions in milk and reconstituted milk replacer should be prepared immediately before use. Any medicated milk or milk replacer which is not consumed within one hour should be discarded.

For treatment in chickens, TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder should be administered in the drinking water to provide 250 or 500 mg/litre apramycin activity for five days. This may be achieved by adding 118.2g TP-Apra 500 WSP Apramycin Sulphate 500mg/g Water Soluble Powder (50g apramycin activity) per 100 to 200 litres of water. Any medicated water which is not consumed within 24 hours should be discarded.

CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to apramycin. Do not use in cats.

WARNINGS & PRECAUTIONS.

- Special precautions for use in animals

Official, national and regional antimicrobial policies should be taken into account when the product is used.

- Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid inhaling the powder while preparing the medicate water.

Avoid contact with skin and eyes. In the event of skin contact, wash thoroughly with soap and water.

In the event of accidental eye contact, wash the affected eye with fresh running water and seek medical attention if irritation persists.

Wash hands after use.

INTERACTIONS WITH OTHER MEDICATIONS

Drinking systems should be clean and free of rust to avoid reduction of activity.

PREGNANCY AND LACTATION

Not intended for use in pregnant or lactating animals. However, laboratory studies in the rat and rabbit have not produced any evidence of a teratogenic effect.

ADVERSE EFFECTS

None known.

OVERDOSE & TREATMENT

Pigs have been given up to nine times the recommended use level in their drinking water for 28 days with no untoward reaction.

Calves were given apramycin in milk replacer daily for five days, at doses up to 120 mg/kg bodyweight (3-times recommended use level). There was no toxic effect.

There was no mortality when chickens were given a single oral dose of 1000 mg/kg bodyweight. Chickens were given up to 5 times the recommended level for 15 days with no untoward reaction.

WITHDRAWAL PERIOD

Pigs: Meat: 14 days

Calves: Meat: 28 days

Chickens: Meat: 7 days

Do not use in laying chickens when eggs are intended for human consumption.

MAXIMUM RESIDUAL LIMIT (MRL)

Cattles

Muscle: 0.5 ppm

Fat: 0.5 ppm

Liver: 5 ppm

Kidney: 15 ppm

Edible offal: 15 ppm

Pigs

Muscle: 0.06 ppm

Fat: 0.06 ppm

Liver: 0.06 ppm

Kidney: 2 ppm

Edible offal: 0.1 ppm

Chicken

Muscle: 0.5 ppm

Fat: 0.5 ppm

Liver: 0.5 ppm

Kidney: 2 ppm

Edible offal: 2 ppm

STORAGE CONDITION

Store in a cool, dry place (below 30°C). Protect from sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.



Product Registration Holder
THYE PHARMA SDN. BHD.

Manufactured by
THYE PHARMA SDN. BHD.

No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised May 2022