

Veterinary Package Insert

TP-TYLOFEN 500MG/G POWDER (50% Tylosin Tartrate)

PRODUCT DESCRIPTION

Off-white to light yellow colour powder.

PHARMACODYNAMICS & PHARMACOKINETICS

Pharmacodynamics:

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

Pharmacokinetics:

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord.

INDICATION

Pigs: For the prevention and control of enzootic pneumonia, and scours caused by organisms (e.g. *Lawsonia intracellularis*) sensitive to tylosin.

Chickens/ Turkeys: For the control of *Mycoplasma synoviae* airsacculitis in chickens and *Mycoplasma gallisepticum* S6 in chickens and turkeys. In the field tylosin has also proved useful in reducing the level of infection following stress associated with live vaccination.

As an aid in the control of outbreaks of necrotic enteritis in chickens caused by *Clostridium perfringens*.

RECOMMENDED DOSAGE AND ADMINISTRATION

It is to be given orally. To be administered in drinking water.

Recommendations for use in chickens and turkeys:

Product is administered in the drinking water at a concentration of 1.1 g per litre.

As an aid in the control of outbreaks of necrotic enteritis caused by *Clostridium perfringens* in chickens use the product in the drinking water for 5 days at a concentration of 0.3 g per litre water (150ppm), to provide 20-50 mg/kg bw, depending on the age and the water consumption of the birds.

Recommendations for use in pigs:

The product is administered in the drinking water to provide 25 mg tylosin/kg bodyweight. This may normally be achieved by adding 0.5 per litre.

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid under-dosing.

Recommended programmes:

a.) Chickens and turkeys

For preventative medication the following programmes are recommended:

As an aid to mycoplasmosis prevention in chickens:

For broiler, layers and replacement pullets that are at 0 to 3 days and 4th week of life, use 76.9g per 1000 birds of product at 1.1g per litre. When layers and replacement pullets on 9-12 week and 16th to 20th week, continue 48 hours medication of product at 1.1g per litre.

For breeders that are at 0 to 3 days, use 109.9g per 1000 birds of product at 1.1g per litre. And at 4th week of life, then use 175.8g per 1000 birds. Then, on 9th, 12th, 16th, 20th and 24th weeks of age, continue 48 hours medication of product at 1.1g per litre.

As an aid to mycoplasmosis prevention in table turkeys that are on 0 to 5 days, use 27.5g per 100 birds of product at 1.1g per litre. Then, on 4th week of life, use 16.5g per 100 birds.

For treatment of mycoplasmosis in broilers, use product at 1.1g per litre for 24 to 72 hours, in layers and replacement pullets, use product at 1.1g per litre for 48 to 72 hours medication and whereas for turkeys, use the same dose for 48 to 120 hours.

As an aid in the control of necrotic enteritis for broilers, layers and replacement pullets, use product at 0.3g per litre for 5 days.

b.) Pigs

A medicated solution of drinking water should generally be administered until 24 hours after scouring or respiratory symptoms have ceased, normally 3-10 days. The diagnosis should be reviewed if there is no response after 5 days of medication.

Product requirements per tonne of pigs daily:

To treat enzootic pneumonia, the product required is 54.9g for approximately 100 litres of water consumption. To treat ileitis, 11 to 22g of product is required for approximately 100 litres of water consumption.

CONTRAINDICATIONS

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or to wildlife.

Do not use in known cases of hypersensitivity to tylosin or other macrolides.

WARNINGS & PRECAUTIONS.

i) Special precautions for use in animals

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the product deviating from the instructions given may increase the prevalence of bacteria resistant to tylosin and may decrease the effectiveness of treatment with other macrolides, lincosamides and streptogramin B due to the potential for cross-resistance.

ii.) Special precautions for the person administering the veterinary medicinal product to animals

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator or a non-disposable respirator with a filter. Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

INTERACTIONS WITH OTHER MEDICATIONS

None known.

PREGNANCY AND LACTATION

No adverse effects to tylosin have been seen in fertility, multi-generation or teratology studies.

ADVERSE EFFECTS

None known.

OVERDOSE & TREATMENT

There is no evidence of tylosin toxicity in animals, at dose rates of up to 1000 mg/kg.

WITHDRAWAL PERIOD

Species	Treatm ent Duratio n (days)	Dose	Meat Withdr awal Period (days)	Egg Withd rawal perio d (days)
Chickens (For the control of chronic respiratory disease)	5	500 mg/litre water (50-200 mg tylosin/kg bw)	1	Zero
Chickens (As an aid in the control of outbreaks of necrotic enteritis caused by <i>Clostridium perfringens</i>)	5	150 mg /litre water (20-50 mg tylosin/kg bw)	Zero	Zero
Turkeys (For the control of chronic respiratory disease)	5	500 mg/litre water	Zero	N/A
Pigs (For the prevention	3-10	250 mg/litre water (25 mg tylosin/kg bw)	Zero	N/A

and control of enzootic pneumonia, swine)				
--	--	--	--	--

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in food µg/kg
Tylosin	Muscle, liver, kidney, skin + fat (Porcine)	100
	Muscle, liver, kidney, skin + fat (Poultry)	100

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use
within 24 hours.

PACKING

500g

1kg

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised April 2017