

EIMERIAX 12% w/w Granular

For Animal Use Only

Active Ingredient:

Salinomycin Sodium

120g/kg

Product Description:

Eimeriax 12% w/w Granular contains Salinomycin mixed with suitable diluents, carriers, and inactive ingredients prepared in a granular form. It is pale brown, homogeneous granule. It is for veterinary use only which aids in prevention of coccidiosis in broiler chickens caused by *Eimeria acervulina*, *Eimeria mitis*, *Eimeria necatrix*, *Eimeria maxima*, *Eimeria tenella* and *Eimeria brunetti*.

Pharmacodynamics:

Salinomycin is a carboxylic polyether ionophore. Its polyether skeleton of the pseudocyclic structure is able to form complexes with metal cations. The lipophilic properties of salinomycin allow it to easily penetrate cellular membranes and disturb the intracellular cation balance which finally leads to the cell death. Owing to these properties, salinomycin shows various biological activities such as growth inhibition of Gram-positive micro-organisms and control of chicken coccidiosis, for which it is applied commercially.

Pharmacokinetics:

Salinomycin is rapidly absorbed, extensively metabolised in chicken and no residues were detected in any of the tissues after 4 days.

Indications:

For the prevention of coccidiosis in broiler chicken caused by *Eimeria acervulina*, *Eimeria mitis*, *Eimeria necatrix*, *Eimeria maxima*, *Eimeria tenella* and *Eimeria brunetti*.

Recommended Dosage:

60 mg/kg (milligrams per kilogram) (0.006%) of Salinomycin Sodium in the complete feed.

Direction for Use: Mix 500g Eimeriax 12% w/w Granular in 1 tonne of finished feed to provide a salinomycin concentration of 60 mg/kg (60ppm). The prepared medicated feed should be fed continuously at the sole ration.

Mode of Administration:

Oral administration

Contraindication:

Due to the chemical properties of the active ingredient, this product is contraindicated for use in dogs, horses and turkeys as it is toxic to these species.

Warning and Precautions:

- Do not feed to horses, dogs, turkeys and any other animal or bird not included on the label.
- Do not feed to dairy cows.
- Do not administer to animals receiving Tiamulin.
- Do not use in birds laying eggs for human consumption.
- Care should be taken when feeding concurrently with other antimicrobial products.
- Operators handling the concentrated product should wear a face mask, gloves and protective clothing.
- Keep this and all drugs out of the reach of children.

Interaction with Other Medicinal Product:

Clinically important, often lethal interactions between the ionophore anticoccidials and the antibiotic tiamulin are a well-known phenomenon in chickens. It has been shown that tiamulin is a potent inhibitor of the activity of hepatic cytochrome P450 enzymes and particularly members of the CYP3A thus delaying

the metabolism of ionophoric polyethers. Subsequently, the latter accumulate following daily ingestions, and signs of toxicity occur due to a relative overdose.

Pregnancy and Lactation:

Not applicable.

Adverse Effects and Toxicity:

None known.

Overdose and Treatment:

No data is available.

Withdrawal Period:

4 days for chicken.

Storage Condition:

Store in a dry cool place at temperatures below 25°C.

Proposed shelf life (after reconstitution):

24 hours.

Dosage Form:

Granules.

Packaging Available:

25 kg in PP bag with polyethylene liner.

Date of Revision of Package Insert: 17th October 2016

Name & Address of Manufacturer:

Inner Mongolia Biok Biology Co., Ltd.

Tuo Electricity Industrial Park, Tuoketuo County, Inner Mongolia, China 010206.

Marketing Authorisation Holder:

Sunzen Biotech Berhad (680889-W)

11, Jalan Anggerik Mokara 31/47, Kota Kemuning, 40460 Shah Alam, Selangor, Malaysia.