

SN-Flubendazole 5%w/w

Oral Powder
(FOR VETERINARY USE ONLY)

COMPOSITION

Flubendazole 5 %w/w

PRODUCT DESCRIPTION

The finished product is a white color powder.

TARGET SPECIES

Pigs

PHARMACODYNAMIC

Flubendazole is a synthetic anthelmintic belonging to the benzimidazole carbamates which acts by inhibiting the microtubular assembly in absorptive cells of nematodes. Flubendazole acts by binding to tubulin, the dimeric subunit protein of the microtubules. It inhibits microtubular assembly in absorptive cells: i.e. of intestinal cells of nematodes. This is shown by disappearance of cytoplasmic microtubules, accumulation of secretory granules in the cytoplasm due to a block in their transport, leading to an impaired coating of the cellular membrane and a decreased digestion and absorption of nutrients. Irreversible lytic degeneration of the cell, due to the accumulation of secretory substances (hydrolytic and proteolytic enzymes), results in the death of the parasite. These changes are relatively fast and are primarily seen in those organelles directly involved in the secretory and absorptive functions of the cells. In contrast the changes are not seen in host cells.

PHARMACOKINETIC

Flubendazole is very poorly soluble in aqueous systems, such as the gastrointestinal tract, which results in a low dissolution rate and a very low absorption. This is reflected by a high faecal excretion of unchanged parent drug. The very small fraction absorbed is extensively metabolised by first-pass metabolism in the liver, involving carbamate hydrolysis and ketone reduction. The biotransformation products are conjugated to glucuronides or sulphate conjugates and excreted in the bile and the urine. The excretion in urine is relatively low and consists almost exclusively of metabolites with only small amounts of unchanged compound. In pigs, highest tissue levels are measured in liver and kidneys. The half-life of flubendazole in tissues is 1 to 2 days.

INDICATIONS

Flubendazole is a broad spectrum anthelmintic for oral administration, active against mature and immature stages of the following nematodes of the gastro-intestinal tract of the pig: *Ascaris suum* (large roundworm), *Hyostrogylus rubidus* (red stomach worm), *Oesophagostomum dentatum* (nodular worm), *Trichuris suis* (whip worm), *Strongyloides ransomi* (threadworm) (adult) & *Metastrongylus apri* (lungworm).

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

None

Special precautions for use:

i. Special precautions for use in animals:

None

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental ingestion by humans should be avoided. May cause sensitisation by skin contact. May cause skin and eye irritation. Avoid direct skin contact. Wear overalls, safety glasses and impervious gloves when mixing and handling the product. Wash affected parts if skin contact occurs. If accidental eye contact occurs, immediately rinse thoroughly with water. If the operations involve potential exposure to dust, wear either a disposable filter and half-mask respirator, or a non-disposable respirator fitted with a filter.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

ADMINISTRATION & DOSAGE

For oral administration only.

For incorporation into dry feed at a registered mill. A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or premixtures containing such products, must be responsible for mixing when incorporation is less than 2 kg per ton for final feed.

Pigs: The standard recommended total dosage is 5 mg flubendazole per kg bodyweight. The amount of product to be incorporated should be calculated according to the average bodyweight of the pigs to be treated.

Add 600 g of product to at least 5 kg of one of the feed ingredients and mix well. Thoroughly mix this premix with the remaining ingredients making in all one tonne of medicated feed, which can then be fed as mash or pellets. This gives 30 mg flubendazole per kg of finished feed. Breeding stock should be treated for 10 consecutive days. Weaners and fattening pigs –should be treated for 5 consecutive days, or the event of a heavy *Trichuris* infestation, for 10 consecutive days.

Treatment frequency: Twice a year unless recommended otherwise by a veterinary surgeon. Pigs brought onto the premises should be treated on arrival and before mixing with other animals.

Treatment of clinical worm infestations at the following intervals:

Lungworm	every 3 - 4 weeks
Nodular worm	every 2 months
Large round worm	every 2 months
Red stomach worm	every month
Whipworm	every 6 weeks

CONTRAINDICATIONS

None known

INTERACTION WITH OTHER MEDICAMENTS

None known

PREGNANCY AND LACTATION

Not applicable

OVERDOSE AND TREATMENT

Flubendazole is an analog of mebendazole for which the side effects of overdose include transient gastrointestinal abnormalities.

SIDE EFFECT

None known

WITHDRAWAL TIME

Animals must not be slaughtered for human consumption during treatment. Pigs may be slaughtered for human consumption only after 7 days from the last treatment.

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store at temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

The powder filled 600 g in aluminum sachets, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 12 April 2019



Product Registration Holder & Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor, Malaysia.